

# KCTD9 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP17624c

#### **Product Information**

Application WB, E Primary Accession Q7L273

Other Accession Q80UN1, NP 060104.2

Reactivity Human **Predicted** Mouse Host Rabbit Clonality Polyclonal Isotype Rabbit IgG **Clone Names** RB37694 **Calculated MW** 42567 179-207 **Antigen Region** 

#### **Additional Information**

**Gene ID** 54793

Other Names BTB/POZ domain-containing protein KCTD9, KCTD9

**Target/Specificity**This KCTD9 antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 179-207 amino acids from the Central

region of human KCTD9.

**Dilution** WB~~1:1000 E~~Use at an assay dependent concentration.

**Format** Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** KCTD9 Antibody (Center) is for research use only and not for use in diagnostic

or therapeutic procedures.

#### **Protein Information**

Name KCTD9

**Function** Substrate-specific adapter of a BCR (BTB-CUL3-RBX1) E3 ubiquitin-protein

ligase complex, which mediates the ubiquitination of target proteins, leading

to their degradation by the proteasome.


## **Background**

KCTD9 contains a potassium channel tetramerisation domain. The N-terminal, cytoplasmic tetramerisation domain (T1) of voltage-gated potassium channels encodes molecular determinants for subfamily-specific assembly of alpha-subunits into functional tetrameric channels. The specific function of KCTD9 is unknown.

### References

Zhou, Y.Y., et al. Zhonghua Gan Zang Bing Za Zhi 16(11):835-839(2008) Lamesch, P., et al. Genomics 89(3):307-315(2007)

## **Images**



KCTD9 Antibody (Center) (Cat. #AP17624c) western blot analysis in Jurkat cell line lysates (35ug/lane). This demonstrates the KCTD9 antibody detected the KCTD9 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.