10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# RNF14 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17648b

## Product Information

| | |
|---|---|
| **Application** | WB, E |
| **Primary Accession** | Q9UBS8 |
| **Other Accession** | NP_899646.1, NP_004281.1 |
| **Reactivity** | Human, Mouse |
| **Predicted** | Rat, Bovine, Canine, Rabbit, Chicken |
| **Host** | Rabbit |
| **Clonality** | Polyclonal |
| **Isotype** | Rabbit IgG |
| **Clone Names** | RB30406 |
| **Calculated MW** | 53837 |
| **Antigen Region** | 336-364 |

## Additional Information

| | |
|---|---|
| **Gene ID** | 9604 |
| **Other Names** | E3 ubiquitin-protein ligase RNF14, 632-, Androgen receptor-associated protein 54, HFB30, RING finger protein 14, Triad2 protein, RNF14, ARA54 |
| **Target/Specificity** | This RNF14 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 336-364 amino acids from the C-terminal region of human RNF14. |
| **Dilution** | WB~~1:1000 E~~Use at an assay dependent concentration. |
| **Format** | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| **Storage** | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| **Precautions** | RNF14 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures. |

## Protein Information

| | |
|---|---|
| **Name** | RNF14 {ECO:0000303\|PubMed:36638793, ECO:0000312\|HGNC:HGNC:10058} |
| **Function** | E3 ubiquitin-protein ligase that plays a key role in the RNF14-RNF25 |

translation quality control pathway, a pathway that takes place when a ribosome has stalled during translation, and which promotes ubiquitination and degradation of translation factors on stalled ribosomes (PubMed:36638793, PubMed:37651229, PubMed:37951215, PubMed:37951216). Recruited to stalled ribosomes by the ribosome collision sensor GCN1 and mediates 'Lys-6'-linked ubiquitination of target proteins, leading to their degradation (PubMed:36638793, PubMed:37651229, PubMed:37951215, PubMed:37951216). Mediates ubiquitination of EEF1A1/eEF1A and ETF1/eRF1 translation factors on stalled ribosomes, leading to their degradation (PubMed:36638793, PubMed:37651229). Also catalyzes ubiquitination of ribosomal proteins RPL0, RPL1, RPL12, RPS13 and RPS17 (PubMed:36638793). Specifically required to resolve RNA-protein cross-links caused by reactive aldehydes, which trigger translation stress by stalling ribosomes: acts by catalying 'Lys-6'-linked ubiquitination of RNA-protein cross-links, leading to their removal by the ATP-dependent unfoldase VCP and subsequent degradation by the proteasome (PubMed:37951215, PubMed:37951216). Independently of its function in the response to stalled ribosomes, acts as a regulator of transcription in Wnt signaling via its interaction with TCF transcription factors (TCF7/TCF1, TCF7L1/TCF3 and TCF7L2/TCF4) (PubMed:23449499). May also play a role as a coactivator for androgen- and, to a lesser extent, progesterone-dependent transcription (PubMed:19345326).

**Cellular Location** Cytoplasm. Nucleus

**Tissue Location** Widely expressed..

## Background


The protein encoded by this gene contains a RING zinc finger, a motif known to be involved in protein-protein interactions. This protein interacts with androgen receptor (AR) and may function as a coactivator that induces AR target gene expression in prostate. A dominant negative mutant of this gene has been demonstrated to inhibit the AR-mediated growth of prostate cancer. This protein also interacts with class III ubiquitin-conjugating enzymes (E2s) and may act as a ubiquitin-ligase (E3) in the ubiquitination of certain nuclear proteins. Five alternatively spliced transcript variants encoding two distinct isoforms have been reported.

## References

Xu, K., et al. Cancer Cell 15(4):270-282(2009)
Lan, K.C., et al. Fertil. Steril. 89 (5 SUPPL), 1397-1405 (2008) :
Kikuchi, H., et al. Carcinogenesis 28(8):1752-1758(2007)
Yang, Z., et al. Endocrinology 148(3):1340-1349(2007)
Yang, Z., et al. Mol. Endocrinol. 21(2):343-358(2007)

## Images

RNF14 Antibody (C-term) (Cat. #AP17648b) western blot analysis in HL-60 cell line lysates (35ug/lane).This demonstrates the RNF14 antibody detected the RNF14 protein (arrow).




Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.