

GTF2A1 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17659b

Product Information

Application	WB, E
Primary Accession	P52655
Other Accession	O08949 , Q99PM3 , NP_963889.1 , NP_056943.1
Reactivity	Human, Mouse
Predicted	Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB36895
Calculated MW	41514
Antigen Region	297-326

Additional Information

Gene ID	2957
Other Names	Transcription initiation factor IIA subunit 1, General transcription factor IIA subunit 1, TFIIAL, Transcription initiation factor TFIIA 42 kDa subunit, TFIIA-42, Transcription initiation factor IIA alpha chain, TFIIA p35 subunit, Transcription initiation factor IIA beta chain, TFIIA p19 subunit, GTF2A1, TF2A1
Target/Specificity	This GTF2A1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 297-326 amino acids from the C-terminal region of human GTF2A1.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	GTF2A1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	GTF2A1
-------------	--------

Synonyms	TF2A1
Function	TFIIA is a component of the transcription machinery of RNA polymerase II and plays an important role in transcriptional activation. TFIIA in a complex with TBP mediates transcriptional activity.
Cellular Location	Nucleus.


Background

Accurate transcription initiation on TATA-containing class II genes involves the ordered assembly of RNA polymerase II (POLR2A; MIM 180660) and several general initiation factors (summarized by DeJong and Roeder, 1993 [PubMed 8224848]). One of these factors is TFIIA, which when purified from HeLa extracts consists of 35-, 19-, and 12-kD subunits.

References

Huang, Y.W., et al. *Oncol. Rep.* 22(4):853-861(2009)
Vega, A., et al. *Gynecol. Oncol.* 112(1):210-214(2009)
Hieb, A.R., et al. *J. Mol. Biol.* 372(3):619-632(2007)
Lee, K.A., et al. *Cytokine* 38(2):96-100(2007)
Ewing, R.M., et al. *Mol. Syst. Biol.* 3, 89 (2007) :

Images



GTF2A1 Antibody (C-term) (Cat. #AP17659b) western blot analysis in mouse kidney tissue lysates (35ug/lane). This demonstrates the GTF2A1 antibody detected the GTF2A1 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.