

# SEPT10 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP17950a

#### **Product Information**

Application WB, E Primary Accession Q9P0V9

Other Accession NP 653311.1, NP 848699.1

Reactivity Human
Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Clone Names RB38137
Calculated MW 52593
Antigen Region 14-40

## **Additional Information**

**Gene ID** 151011

Other Names Septin-10, SEPT10

Target/Specificity This SEPT10 antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 14-40 amino acids from the N-terminal

region of human SEPT10.

**Dilution** WB~~1:1000 E~~Use at an assay dependent concentration.

**Format** Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

**Storage** Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** SEPT10 Antibody (N-term) is for research use only and not for use in

diagnostic or therapeutic procedures.

#### **Protein Information**

Name SEPTIN10 ( HGNC:14349)

Synonyms SEPT10

**Function** Filament-forming cytoskeletal GTPase. May play a role in cytokinesis

(Potential).

**Cellular Location** Cytoplasm. Cytoplasm, cytoskeleton. Cell projection, cilium, flagellum

{ECO:0000250 | UniProtKB:Q8C650}. Note=Detected in the annulus of the sperm flagellum and in the neck region in spermatids and mature sperm (By

similarity). Using a GFP-fusion protein, detected in the nucleus.

{ECO:0000250|UniProtKB:Q8C650, ECO:0000269|PubMed:12711328}

**Tissue Location** Widely expressed. Abundantly expressed in heart and kidney, placenta,

skeletal muscles, liver and lung, as well as various tumor cell lines.


# **Background**

This gene encodes a member of the septin family of cytoskeletal proteins with GTPase activity. This protein localizes to the cytoplasm and nucleus and displays GTP-binding and GTPase activity. Alternate splicing results in two transcript variants encoding different isoforms.

### References

Benedetti, D., et al. Leukemia 22(1):224-226(2008) Lamesch, P., et al. Genomics 89(3):307-315(2007) Hillier, L.W., et al. Nature 434(7034):724-731(2005) Sui, L., et al. Biochem. Biophys. Res. Commun. 304(2):393-398(2003)

# **Images**



SEPT10 Antibody (N-term) (Cat. #AP17950a) western blot analysis in MDA-MB231 cell line lysates (35ug/lane). This demonstrates the SEPT10 antibody detected the SEPT10 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.