

# SNAP25 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP18126C

## Product Information

---

<b>Application</b>	WB, E
<b>Primary Accession</b>	<a href="#">P60880</a>
<b>Other Accession</b>	<a href="#">P60881</a> , <a href="#">P60879</a> , <a href="#">P60878</a> , <a href="#">Q17QQ3</a> , <a href="#">Q6PC54</a> , <a href="#">Q5TZ66</a> , <a href="#">NP_003072.2</a>
<b>Reactivity</b>	Human
<b>Predicted</b>	Zebrafish, Bovine, Chicken, Mouse, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB21262
<b>Calculated MW</b>	23315
<b>Antigen Region</b>	36-63

## Additional Information

---

<b>Gene ID</b>	6616
<b>Other Names</b>	Synaptosomal-associated protein 25, SNAP-25, Super protein, SUP, Synaptosomal-associated 25 kDa protein, SNAP25, SNAP
<b>Target/Specificity</b>	This SNAP25 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 36-63 amino acids from the Central region of human SNAP25.
<b>Dilution</b>	WB~~1:1000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	SNAP25 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	SNAP25
<b>Synonyms</b>	SNAP

<b>Function</b>	t-SNARE involved in the molecular regulation of neurotransmitter release. May play an important role in the synaptic function of specific neuronal systems. Associates with proteins involved in vesicle docking and membrane fusion. Regulates plasma membrane recycling through its interaction with CENPF. Modulates the gating characteristics of the delayed rectifier voltage-dependent potassium channel KCNB1 in pancreatic beta cells.
<b>Cellular Location</b>	Cytoplasm, perinuclear region {ECO:0000250 UniProtKB:P60879}. Cell membrane {ECO:0000250 UniProtKB:P60881}; Lipid-anchor {ECO:0000250 UniProtKB:P60879}. Synapse, synaptosome {ECO:0000250 UniProtKB:P60879}. Photoreceptor inner segment {ECO:0000250 UniProtKB:P60879}. Note=Membrane association requires palmitoylation. Expressed throughout cytoplasm, concentrating at the perinuclear region. Colocalizes with KCNB1 at the cell membrane (By similarity). Colocalizes with PLCL1 at the cell membrane (By similarity). {ECO:0000250 UniProtKB:P60879, ECO:0000250 UniProtKB:P60881}
<b>Tissue Location</b>	Neurons of the neocortex, hippocampus, piriform cortex, anterior thalamic nuclei, pontine nuclei, and granule cells of the cerebellum


## Background

Synaptic vesicle membrane docking and fusion is mediated by SNAREs (soluble N-ethylmaleimide-sensitive factor attachment protein receptors) located on the vesicle membrane (v-SNAREs) and the target membrane (t-SNAREs). The assembled v-SNARE/t-SNARE complex consists of a bundle of four helices, one of which is supplied by v-SNARE and the other three by t-SNARE. For t-SNAREs on the plasma membrane, the protein syntaxin supplies one helix and the protein encoded by this gene contributes the other two. Therefore, this gene product is a presynaptic plasma membrane protein involved in the regulation of neurotransmitter release. Two alternative transcript variants encoding different protein isoforms have been described for this gene.

## References


Shimada, M., et al. Hum. Genet. 128(4):433-441(2010)  
Greaves, J., et al. J. Biol. Chem. 285(32):24629-24638(2010)  
Oner, O., et al. J Atten Disord (2010) In press :  
Zhang, H., et al. Eur. J. Paediatr. Neurol. (2010) In press :  
Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :

## Images



Anti-SNAP25 Antibody (Center) at 1:1000 dilution + SH-SY5Y whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 23 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

SNAP25 Antibody (Center) (Cat. #AP18126c) western blot analysis in 293 cell line lysates (35ug/lane).This



demonstrates the SNAP25 antibody detected the SNAP25 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.