10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

OR4S2 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP18191b

Product Information

Application	WB, E
Primary Accession	Q8NH73
Other Accession	NP_001004059.2
Reactivity	Human
Predicted	Mouse, Rat, Bovine, Canine, Rabbit
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB28918
Calculated MW	35172
Antigen Region	206-233

Additional Information

Gene ID	219431
Other Names	Olfactory receptor 4S2, Olfactory receptor OR11-137, OR4S2, OR4S2P
Target/Specificity	This OR4S2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 206-233 amino acids from the C-terminal region of human OR4S2.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	OR4S2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	OR4S2
Synonyms	OR4S2P
Function	Odorant receptor.

Cellular Location Cell membrane; Multi-pass membrane protein.


Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

References

Malnic, B., et al. Proc. Natl. Acad. Sci. U.S.A. 101(8):2584-2589(2004)
Gilad, Y., et al. Mol. Biol. Evol. 20(3):307-314(2003)
Fuchs, T., et al. Genomics 80(3):295-302(2002)

Images



OR4S2 Antibody (C-term) (Cat. #AP18191b) western blot analysis in K562 cell line lysates (35ug/lane).This demonstrates the OR4S2 antibody detected the OR4S2 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.