

TCEAL2 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP18289a

Product Information

Application	WB, E
Primary Accession	Q9H3H9
Other Accession	NP_525129.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB38366
Calculated MW	25850
Antigen Region	1-30

Additional Information

Gene ID	140597
Other Names	Transcription elongation factor A protein-like 2, TCEA-like protein 2, Transcription elongation factor S-II protein-like 2, TCEAL2
Target/Specificity	This TCEAL2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human TCEAL2.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	TCEAL2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TCEAL2
Function	May be involved in transcriptional regulation.
Cellular Location	Nucleus.


Background

This gene encodes a member of the transcription elongation factor A (SII)-like (TCEAL) gene family. Members of this family contain TFA domains and may function as nuclear phosphoproteins that modulate transcription in a promoter context-dependent manner. Multiple family members are located on the X chromosome. [provided by RefSeq].

References

Ross, M.T., et al. Nature 434(7031):325-337(2005)
Pillutla, R.C., et al. Genomics 56(2):217-220(1999)
Bussow, K., et al. Nucleic Acids Res. 26(21):5007-5008(1998)

Images



Western blot analysis of lysate from HepG2 cell line, using TCEAL2 Antibody (N-term)(Cat. #AP18289a). AP18289a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug per lane.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.