

TACO1 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP18395a

Product Information

Application WB, E **Primary Accession** Q9BSH4 Other Accession NP 057444.2 Reactivity Human Host Rabbit Clonality Polyclonal Isotype Rabbit IgG **Clone Names** RB38422 **Calculated MW** 32477 50-77 **Antigen Region**

Additional Information

Gene ID 51204

Other Names Translational activator of cytochrome c oxidase 1, Coiled-coil

domain-containing protein 44, Translational activator of

mitochondrially-encoded cytochrome c oxidase I, TACO1, CCDC44

Target/Specificity This TACO1 antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 50-77 amino acids from the N-terminal

region of human TACO1.

Dilution WB~~1:1000 E~~Use at an assay dependent concentration.

Format Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions TACO1 Antibody (N-term) is for research use only and not for use in

diagnostic or therapeutic procedures.

Protein Information

Name TACO1

Synonyms CCDC44

Function Acts as a translational activator of mitochondrially-encoded cytochrome c

oxidase 1.

Cellular Location Mitochondrion.


Background

This gene encodes a mitochondrial protein that function as a translational activator of mitochondrially-encoded cytochrome c oxidase 1. Mutations in this gene are associated with Leigh syndrome.

References

Weraarpachai, W., et al. Nat. Genet. 41(7):833-837(2009) Lamesch, P., et al. Genomics 89(3):307-315(2007)

Images



TACO1 Antibody (N-term) (Cat. #AP18395a) western blot analysis in 293 cell line lysates (35ug/lane). This demonstrates the TACO1 Antibody detected the TACO1 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.