10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

TRHR Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP18399c

Product Information

Application	WB, E
Primary Accession	P34981
Other Accession	Q01717, NP_003292.1
Reactivity	Human
Predicted	Rat, Mouse, Bovine, Canine, Rabbit, Chicken
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB38442
Calculated MW	45085
Antigen Region	209-238

Additional Information

Gene ID	7201
Other Names	Thyrotropin-releasing hormone receptor, TRH-R, Thyroliberin receptor, TRHR
Target/Specificity	This TRHR antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 209-238 amino acids from the Central region of human TRHR.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	TRHR Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TRHR
Function	Receptor for thyrotropin-releasing hormone (TRH). Upon ligand binding, this G protein-coupled receptor triggers activation of the phosphatidylinositol (IP3)-calcium-protein kinase C (PKC) pathway.

Cellular Location Cell membrane; Multi-pass membrane protein


Background

Thyrotropin-releasing hormone (TRH; MIM 275120), a small neuropeptide, is widely distributed throughout the central and peripheral nervous system as well as in extraneural tissues. The peptide is synthesized in the hypothalamus and transported by the portal vascular system to the anterior pituitary where it acts on thyrotropic and lactotropic cells to promote secretion of TSH and prolactin, respectively. Thyrotropin-releasing hormone receptor is a G protein-coupled receptor that activates the inositol phospholipid-calcium-protein kinase C transduction pathway upon the binding of TRH. The TRHR gene is expressed in the thyrotrope cells of the anterior pituitary.

References

Gehret, A.U., et al. Biochem. J. 428(2):235-245(2010)
Gehret, A.U., et al. Mol. Pharmacol. 77(2):288-297(2010)
Zhu, C.C., et al. J. Biol. Chem. 277(31):28228-28237(2002)
Matre, V., et al. J. Neurochem. 72(1):40-50(1999)
Iwasaki, T., et al. J. Biol. Chem. 271(36):22183-22188(1996)

Images



TRHR Antibody (Center) (Cat. #AP18399c) western blot analysis in Jurkat cell line lysates (35ug/lane).This demonstrates the TRHR Antibody detected the TRHR protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.