

# UVRAG Antibody (L133)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP1850d

## Product Information

---

<b>Application</b>	WB, IF, IHC-P, E
<b>Primary Accession</b>	<a href="#">Q9P2Y5</a>
<b>Reactivity</b>	Human, Mouse
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Names</b>	RB10846
<b>Calculated MW</b>	78151
<b>Antigen Region</b>	120-152

## Additional Information

---

<b>Gene ID</b>	7405
<b>Other Names</b>	UV radiation resistance-associated gene protein, p63, UVRAG
<b>Target/Specificity</b>	This UVRAG antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 120-152 amino acids from human UVRAG.
<b>Dilution</b>	WB~~1:1000 IF~~Tested IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	UVRAG Antibody (L133) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	UVRAG
<b>Function</b>	Versatile protein that is involved in regulation of different cellular pathways implicated in membrane trafficking. Involved in regulation of the COPI-dependent retrograde transport from Golgi and the endoplasmic reticulum by associating with the NRZ complex; the function is dependent on

its binding to phosphatidylinositol 3- phosphate (PtdIns(3)P) (PubMed:[16799551](#), PubMed:[18552835](#), PubMed:[20643123](#), PubMed:[24056303](#), PubMed:[28306502](#)). During autophagy acts as a regulatory subunit of the alternative PI3K complex II (PI3KC3-C2) that mediates formation of phosphatidylinositol 3-phosphate and is believed to be involved in maturation of autophagosomes and endocytosis. Activates lipid kinase activity of PIK3C3 (PubMed:[16799551](#), PubMed:[20643123](#), PubMed:[24056303](#), PubMed:[28306502](#)). Involved in the regulation of degradative endocytic trafficking and cytokinesis, and in regulation of ATG9A transport from the Golgi to the autophagosome; the functions seems to implicate its association with PI3KC3-C2 (PubMed:[16799551](#), PubMed:[20643123](#), PubMed:[24056303](#)). Involved in maturation of autophagosomes and degradative endocytic trafficking independently of BECN1 but depending on its association with a class C Vps complex (possibly the HOPS complex); the association is also proposed to promote autophagosome recruitment and activation of Rab7 and endosome-endosome fusion events (PubMed:[18552835](#), PubMed:[28306502](#)). Enhances class C Vps complex (possibly HOPS complex) association with a SNARE complex and promotes fusogenic SNARE complex formation during late endocytic membrane fusion (PubMed:[24550300](#)). In case of negative-strand RNA virus infection is required for efficient virus entry, promotes endocytic transport of virions and is implicated in a VAMP8- specific fusogenic SNARE complex assembly (PubMed:[24550300](#)).

#### Cellular Location

Late endosome. Lysosome. Cytoplasmic vesicle, autophagosome. Early endosome. Endoplasmic reticulum. Midbody. Chromosome, centromere. Note=Colocalizes with RAB9-positive compartments involved in retrograde transport from late endosomes to trans-Golgi network. Colocalization with early endosomes is only partial (PubMed:[24056303](#)). Recruited to autophagosome following interaction with RUBCNL/PACER (PubMed:[28306502](#))

#### Tissue Location

Highly expressed in brain, lung, kidney and liver.

## Background


UVRAG complements the ultraviolet sensitivity of xeroderma pigmentosum group C cells and encodes a protein with a C2 domain. The protein activates the Beclin1-PI(3)KC3 complex, promoting autophagy and suppressing the proliferation and tumorigenicity of human colon cancer cells. Chromosomal aberrations involving this gene are associated with left-right axis malformation and mutations in this gene have been associated with colon cancer.

## References


- Liang,C., et al. Nat. Cell Biol. 8 (7), 688-699 (2006)  
 Ionov,Y., et al. Oncogene 23 (3), 639-645 (2004)  
 Goi,T., et al. Surg. Today 33 (9), 702-706 (2003)  
 Iida,A., et al. Hum. Genet. 106 (3), 277-287 (2000)  
 Perelman,B., et al. Genomics 41 (3), 397-405 (1997)  
 Bekri,S., et al. Cytogenet. Cell Genet. 79 (1-2), 125-131 (1997)  
 Teitz,T., et al. Gene 87 (2), 295-298 (1990)

## Images


Western blot analysis of anti-hUVRAG (Center) Pab (RB10846) in mouse liver tissue lysate. hUVRAG



(Center)(arrow) was detected using the purified Pab.



Immunofluorescence staining of Autophagy UVRAG antibody (Cat# AP1850d) on Methanol-fixed HeLa cells. Data courtesy of Dr. Eeva-Liisa Eskelinen, University of Helsinki, Finland.



Formalin-fixed and paraffin-embedded human hepatocarcinoma tissue reacted with hUVRAG (N-term L133), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.