

GOLGA8B Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP18554a

Product Information

Application	WB, E
Primary Accession	A8MQT2
Other Accession	NP_001018861.3
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB38798
Calculated MW	67278
Antigen Region	18-44

Additional Information

Gene ID	440270
Other Names	Golgin subfamily A member 8B, Golgin-67, GOLGA8B
Target/Specificity	This GOLGA8B antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 18-44 amino acids from the N-terminal region of human GOLGA8B.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	GOLGA8B Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	GOLGA8B
Function	May be involved in maintaining Golgi structure.
Cellular Location	Golgi apparatus, Golgi stack membrane; Peripheral membrane protein

Tissue Location

Highly expressed in brain, heart and kidney. Detected at lower levels in liver, thymus, spleen, lung and peripheral blood leukocytes.

Background

GOLGA8B may be involved in maintaining Golgi structure (By similarity).


References

Zody, M.C., et al. Nature 440(7084):671-675(2006)

Eystathiou, T., et al. J. Autoimmun. 14(2):179-187(2000)

Jakymiw, A., et al. J. Biol. Chem. 275(6):4137-4144(2000)

Images



All lanes: Anti-GOLGA8B Antibody (N-term) at 1:2000 dilution Lane 1: CCRF-CEM whole cell lysate Lane 2: HeLa whole cell lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 67 KDa Blocking/Dilution buffer: 5% NFDm/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.