

Mouse Il17ra Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP18687b

Product Information

Application	WB, E
Primary Accession	Q60943
Other Accession	NP_032385.1
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB38645
Calculated MW	97808
Antigen Region	721-747

Additional Information

Gene ID	16172
Other Names	Interleukin-17 receptor A, IL-17 receptor A, IL-17RA, CD217, Il17ra, Il17r
Target/Specificity	This Mouse Il17ra antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 721-747 amino acids of mouse Il17ra.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Mouse Il17ra Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	Il17ra
Synonyms	Il17r
Function	Receptor for IL17A and IL17F, major effector cytokines of innate and adaptive immune system involved in antimicrobial host defense and maintenance of tissue integrity. Receptor for IL17A (PubMed: 17911633 ,

PubMed:[20554964](#), PubMed:[27923703](#), PubMed:[8777726](#)). Receptor for IL17F (PubMed:[17911633](#), PubMed:[20554964](#)). Binds to IL17A with higher affinity than to IL17F (PubMed:[17911633](#)). Binds IL17A and IL17F homodimers as part of a heterodimeric complex with IL17RC (By similarity). Also binds heterodimers formed by IL17A and IL17F as part of a heterodimeric complex with IL17RC (By similarity). Cytokine binding triggers homotypic interaction of IL17RA and IL17RC chains with TRAF3IP2 adapter, leading to TRAF6-mediated activation of NF-kappa-B and MAPkinase pathways, ultimately resulting in transcriptional activation of cytokines, chemokines, antimicrobial peptides and matrix metalloproteinases, with potential strong immune inflammation (By similarity). Involved in antimicrobial host defense primarily promoting neutrophil activation and recruitment at infection sites to destroy extracellular bacteria and fungi (PubMed:[20364087](#), PubMed:[21993848](#)). In secondary lymphoid organs, contributes to germinal center formation by regulating the chemotactic response of B cells to CXCL12 and CXCL13, enhancing retention of B cells within the germinal centers, B cell somatic hypermutation rate and selection toward plasma cells (PubMed:[18157131](#)). Plays a role in the maintenance of the integrity of epithelial barriers during homeostasis and pathogen infection. Stimulates the production of antimicrobial beta-defensins DEFB1, DEFB103A, and DEFB104A by mucosal epithelial cells, limiting the entry of microbes through the epithelial barriers (PubMed:[19144317](#)). Involved in antiviral host defense through various mechanisms. Enhances immunity against West Nile virus by promoting T cell cytotoxicity (PubMed:[27795421](#)). Contributes to influenza A virus (H1N1) clearance by driving the differentiation of B-1a B cells, providing for production of virus-specific IgM antibodies at first line of host defense (PubMed:[26735852](#)). Receptor for IL17C as part of a heterodimeric complex with IL17RE (PubMed:[21982598](#), PubMed:[21993848](#), PubMed:[21993849](#)).

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location

Widely expressed (PubMed:[21993848](#)). Highly expressed in T cells and macrophages (PubMed:[19144317](#)). Highly expressed in B-1a B cells and at a lower extent in B-1b and B-2 B cells (at protein level) (PubMed:[26735852](#)).

Background


Receptor for IL17A. Binds its ligand with low affinity, suggesting that additional components are involved in IL17A-induced signaling (By similarity).

References

- Onishi, R.M., et al. J. Biol. Chem. 285(43):32751-32759(2010)
 Guiton, R., et al. J. Infect. Dis. 202(3):427-435(2010)
 Mitsdoerffer, M., et al. Proc. Natl. Acad. Sci. U.S.A. 107(32):14292-14297(2010)
 Hill, G.R., et al. Blood 116(5):819-828(2010)
 Wu, H.J., et al. Immunity 32(6):815-827(2010)

Images

Mouse Il17ra Antibody (C-term) (Cat. #AP18687b)
 western blot analysis in mouse NIH-3T3 cell line lysates
 (35ug/lane). This demonstrates the Mouse Il17ra antibody
 detected the Mouse Il17ra protein (arrow).



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.