

IRG1 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP18784c

Product Information

Application	WB, E
Primary Accession	A6NK06
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB38857
Calculated MW	52628
Antigen Region	306-332

Additional Information

Gene ID	730249
Other Names	Cis-aconitate decarboxylase, CAD, Aconitate decarboxylase, Cis-aconitic acid decarboxylase, Immune-responsive gene 1 protein, IRG1
Target/Specificity	This IRG1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 306-332 amino acids from the Central region of human IRG1.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	IRG1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ACOD1 (HGNC:33904)
Function	Cis-aconitate decarboxylase that catalyzes production of itaconate and is involved in the inhibition of the inflammatory response (PubMed: 23609450 , PubMed: 23610393 , PubMed: 31548418 , PubMed: 35662396). Acts as a negative regulator of the Toll-like receptors (TLRs)-mediated inflammatory innate

response by stimulating the tumor necrosis factor alpha-induced protein TNFAIP3 expression via reactive oxygen species (ROS) in LPS-tolerized macrophages (PubMed:[23609450](#)). Involved in antimicrobial response of innate immune cells; ACOD1-mediated itaconic acid production contributes to the antimicrobial activity of macrophages by generating itaconate, leading to alkylation of proteins, such as TFEB (PubMed:[23610393](#), PubMed:[35662396](#)). Involved in antiviral response following infection by flavivirus in neurons: ACOD1-mediated itaconate production inhibits the activity of succinate dehydrogenase, generating a metabolic state in neurons that suppresses replication of viral genomes (By similarity). Plays a role in the embryo implantation (By similarity).

Cellular Location

Mitochondrion {ECO:0000250 | UniProtKB:P54987}.


Tissue Location

Expressed in LPS-tolerized macrophages (at protein level). Expressed in peripheral blood mononuclear cells (PBMCs), microglia and macrophage cells.

Background

The function of this protein remains unknown.

Images



IRG1 Antibody (Center)(Cat. #AP18784c) western blot analysis in HepG2 cell line lysates (35ug/lane). This demonstrates the IRG1 antibody detected the IRG1 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.