

POLR1A Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP18976b

Product Information

Application	WB, E
Primary Accession	O95602
Other Accession	NP_056240.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB39202
Calculated MW	194811
Antigen Region	1437-1465

Additional Information

Gene ID	25885
Other Names	DNA-directed RNA polymerase I subunit RPA1, RNA polymerase I subunit A1, A190, DNA-directed RNA polymerase I largest subunit, DNA-directed RNA polymerase I subunit A, RNA polymerase I 194 kDa subunit, RPA194, POLR1A
Target/Specificity	This POLR1A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1437-1465 amino acids from the C-terminal region of human POLR1A.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	POLR1A Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	POLR1A {ECO:0000303 PubMed:25913037, ECO:0000312 HGNC:HGNC:17264}
Function	Catalytic core component of RNA polymerase I (Pol I), a DNA- dependent

RNA polymerase which synthesizes ribosomal RNA precursors using the four ribonucleoside triphosphates as substrates. Transcribes 47S pre-rRNAs from multicopy rRNA gene clusters, giving rise to 5.8S, 18S and 28S ribosomal RNAs (PubMed:[11250903](#), PubMed:[11283244](#), PubMed:[16858408](#), PubMed:[34671025](#), PubMed:[34887565](#), PubMed:[36271492](#)). Pol I-mediated transcription cycle proceeds through transcription initiation, transcription elongation and transcription termination stages. During transcription initiation, Pol I pre-initiation complex (PIC) is recruited by the selectivity factor 1 (SL1/TIF-IB) complex bound to the core promoter that precedes an rDNA repeat unit. The PIC assembly bends the promoter favoring the formation of the transcription bubble and promoter escape. Once the polymerase has escaped from the promoter it enters the elongation phase during which RNA is actively polymerized, based on complementarity with the template DNA strand. Highly processive, assembles in structures referred to as 'Miller trees' where many elongating Pol I complexes queue and transcribe the same rDNA coding regions. At terminator sequences downstream of the rDNA gene, PTRF interacts with Pol I and halts Pol I transcription leading to the release of the RNA transcript and polymerase from the DNA (PubMed:[11250903](#), PubMed:[11283244](#), PubMed:[16858408](#), PubMed:[34671025](#), PubMed:[34887565](#), PubMed:[36271492](#)). Forms Pol I active center together with the second largest subunit POLR1B/RPA2. Appends one nucleotide at a time to the 3' end of the nascent RNA, with POLR1A/RPA1 contributing a Mg(2+)-coordinating DxGDG motif, and POLR1B/RPA2 participating in the coordination of a second Mg(2+) ion and providing lysine residues believed to facilitate Watson-Crick base pairing between the incoming nucleotide and the template base. Typically, Mg(2+) ions direct a 5' nucleoside triphosphate to form a phosphodiester bond with the 3' hydroxyl of the preceding nucleotide of the nascent RNA, with the elimination of pyrophosphate. Has proofreading activity: Pauses and backtracks to allow the cleavage of a missincorporated nucleotide via POLR1H/RPA12. High Pol I processivity is associated with decreased transcription fidelity (By similarity) (PubMed:[11250903](#), PubMed:[11283244](#), PubMed:[16858408](#), PubMed:[34671025](#), PubMed:[34887565](#), PubMed:[36271492](#)).

Cellular Location

Nucleus, nucleolus. Chromosome {ECO:0000250|UniProtKB:O35134}

Background


DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA using the four ribonucleoside triphosphates as substrates. Largest and catalytic core component of RNA polymerase I which synthesizes ribosomal RNA precursors. Forms the polymerase active center together with the second largest subunit. A single stranded DNA template strand of the promoter is positioned within the central active site cleft of Pol I. A bridging helix emanates from RPA1 and crosses the cleft near the catalytic site and is thought to promote translocation of Pol I by acting as a ratchet that moves the RNA-DNA hybrid through the active site by switching from straight to bent conformations at each step of nucleotide addition (By similarity).

References

Rose, J. Phd, et al. Mol. Med. (2010) In press : Percipalle, P., et al. EMBO Rep. 7(5):525-530(2006)
 Bouwmeester, T., et al. Nat. Cell Biol. 6(2):97-105(2004) Hirschler-Laszkiewicz, I., et al. J. Biol. Chem. 278(21):18953-18959(2003) Dundr, M., et al. Science 298(5598):1623-1626(2002)

Images

POLR1A Antibody (C-term) (Cat. #AP18976b) western blot analysis in A375 cell line lysates (35ug/lane).This



demonstrates the POLR1A antibody detected the POLR1A protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.