

# Mouse Gsta3 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP19311b

## Product Information

---

Application	WB, E
Primary Accession	<a href="#">P30115</a>
Other Accession	<a href="#">NP_001070821.1</a>
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB40186
Calculated MW	25361
Antigen Region	183-211

## Additional Information

---

Gene ID	14859
Other Names	Glutathione S-transferase A3, GST class-alpha member 3, Glutathione S-transferase Ya3, Glutathione S-transferase Yc, Gsta3, Gstyc
Target/Specificity	This Mouse Gsta3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 183-211 amino acids from the C-terminal region of mouse Gsta3.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Mouse Gsta3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	Gsta3 {ECO:0000312 MGI:MGI:95856}
Synonyms	Gstyc
Function	Conjugation of reduced glutathione to a wide number of exogenous and

endogenous hydrophobic electrophiles. Catalyzes isomerization reactions that contribute to the biosynthesis of steroid hormones. Efficiently catalyze obligatory double-bond isomerizations of delta(5)-androstene-3,17-dione and delta(5)-pregnene-3,20-dione, precursors to testosterone and progesterone, respectively (By similarity). Has a high catalytic activity for aflatoxin B1-8,9 epoxide (PubMed:[1637297](#)).

**Cellular Location** Cytoplasm.

## Background

---

Conjugation of reduced glutathione to a wide number of exogenous and endogenous hydrophobic electrophiles. This GST has a high catalytic activity for aflatoxin B1-8,9 epoxide.


## References

---

Ilic, Z., et al. Toxicol. Appl. Pharmacol. 242(3):241-246(2010)  
Caruana, G., et al. Gene Expr. Patterns 6(8):807-825(2006)  
Brown, A.C., et al. BMC Genet. 6, 12 (2005) :  
Wang, Y., et al. Cancer Res. 64(5):1647-1654(2004)  
Jowsey, I.R., et al. Biochem. Biophys. Res. Commun. 312(4):1226-1235(2003)

## Images

---



Anti-Mouse Gsta3 Antibody (C-term) at 1:4000 dilution + mouse bladder lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 25 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.