

# EIF3E Antibody(Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP19409c

## **Product Information**

Application	WB, E
Primary Accession	<u>P60228</u>
Other Accession	<u>Q641X8, P60229, Q4R6G8, Q5ZLA5, Q3T102, Q3B8M3, Q1LUA8, Q05AY2,</u>
	<u>Q6DRI1, NP_001559.1</u>
Reactivity	Human, Zebrafish
Predicted	Zebrafish, Xenopus, Bovine, Chicken, Monkey, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	52221
Antigen Region	248-276

## **Additional Information**

Gene ID	3646
Other Names	Eukaryotic translation initiation factor 3 subunit E {ECO:0000255 HAMAP-Rule:MF_03004}, eIF3e {ECO:0000255 HAMAP-Rule:MF_03004}, Eukaryotic translation initiation factor 3 subunit 6 {ECO:0000255 HAMAP-Rule:MF_03004}, Viral integration site protein INT-6 homolog, eIF-3 p48 {ECO:0000255 HAMAP-Rule:MF_03004}, EIF3E {ECO:0000255 HAMAP-Rule:MF_03004}
Target/Specificity	This EIF3E antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 248-276 amino acids from the Central region of human EIF3E.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	EIF3E Antibody(Center) is for research use only and not for use in diagnostic or therapeutic procedures.

# **Protein Information**

Name	EIF3E {ECO:0000255 HAMAP-Rule:MF_03004}
Function	Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis (PubMed: <u>17581632</u> , PubMed: <u>25849773</u> , PubMed: <u>27462815</u> ). The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl- tRNAi and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation (PubMed: <u>17581632</u> ). The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression (PubMed: <u>25849773</u> ). Required for nonsense-mediated mRNA decay (NMD); may act in conjunction with UPF2 to divert mRNAs from translation to the NMD pathway (PubMed: <u>17468741</u> ). May interact with MCM7 and EPAS1 and regulate the proteasome-mediated degradation of these proteins (PubMed: <u>17310990</u> , PubMed: <u>17324924</u> ).
Cellular Location	Cytoplasm. Nucleus, PML body.
Tissue Location	Ubiquitously expressed. Expressed at highest levels in appendix, lymph, pancreas, skeletal muscle, spleen and thymus

#### Background

Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNAi and eIF-5 to form the 43S preinitiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of posttermination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. Required for nonsense-mediated mRNA decay (NMD); may act in conjunction with UPF2 to divert mRNAs from translation to the NMD pathway. May interact with MCM7 and EPAS1 and regulate the proteasome-mediated degradation of these proteins.

## References

Grzmil, M., et al. Oncogene 29(28):4080-4089(2010) Zhou, M., et al. Proc. Natl. Acad. Sci. U.S.A. 105(47):18139-18144(2008) Masutani, M., et al. EMBO J. 26(14):3373-3383(2007) Morris, C., et al. EMBO Rep. 8(6):596-602(2007) Sirchia, R., et al. Biol. Chem. 388(5):457-465(2007)

#### Images

DANRE eif3eb Antibody (Center) (Cat. #AP19409c) western blot analysis in zebra fish brain tissue lysates (35ug/lane).This demonstrates the DANRE eif3eb antibody detected the DANRE eif3eb protein (arrow).



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.