10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

ASCL1 Antibody(N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP19731a

Product Information

Application	WB, E
Primary Accession	P50553
Other Accession	NP_004307.2
Reactivity	Human, Rat, Mouse
Predicted	Bovine, Canine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	25454
Antigen Region	63-90

Additional Information

Gene ID	429
Other Names	Achaete-scute homolog 1, ASH-1, hASH1, Class A basic helix-loop-helix protein 46, bHLHa46, ASCL1, ASH1, BHLHA46, HASH1
Target/Specificity	This ASCL1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 63-90 amino acids from the N-terminal region of human ASCL1.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ASCL1 Antibody(N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ASCL1 (HGNC:738)
Function	Transcription factor that plays a key role in neuronal differentiation: acts as a pioneer transcription factor, accessing closed chromatin to allow other factors to bind and activate neural pathways. Directly binds the E box motif

(5'-CANNTG-3') on promoters and promotes transcription of neuronal genes. The combination of three transcription factors, ASCL1, POU3F2/BRN2 and MYT1L, is sufficient to reprogram fibroblasts and other somatic cells into induced neuronal (iN) cells in vitro. Plays a role at early stages of development of specific neural lineages in most regions of the CNS, and of several lineages in the PNS. Essential for the generation of olfactory and autonomic neurons. Acts synergistically with FOXN4 to specify the identity of V2b neurons rather than V2a from bipotential p2 progenitors during spinal cord neurogenesis, probably through DLL4-NOTCH signaling activation. Involved in the regulation of neuroendocrine cell development in the glandular stomach (By similarity).

Cellular Location Nucleus {ECO:0000250|UniProtKB:Q02067}.


Background

This gene encodes a member of the basic helix-loop-helix (BHLH) family of transcription factors. The protein activates transcription by binding to the E box (5'-CANNTG-3'). Dimerization with other BHLH proteins is required for efficient DNA binding. This protein plays a role in the neuronal commitment and differentiation and in the generation of olfactory and autonomic neurons. Mutations in this gene may contribute to the congenital central hypoventilation syndrome (CCHS) phenotype in rare cases.

References

Phi, J.H., et al. J Neurosurg Pediatr 5(6):608-614(2010)
McGaughey, D.M., et al. Genomics 95(6):363-369(2010)
Li, F., et al. Zhongguo Fei Ai Za Zhi 13(4):317-321(2010)
Deng, H., et al. Biochem. Biophys. Res. Commun. 392(4):548-550(2010)
Improgo, M.R., et al. Mol. Cancer Res. 8(2):194-203(2010)

Images



ASCL1 Antibody (N-term) (Cat. #AP19731a) western blot analysis in NCI-H460 cell line lysates (35ug/lane).This demonstrates the ASCL1 antibody detected the ASCL1 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.