

NR2E1 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP19904c

Product Information

Application	WB, E
Primary Accession	Q9Y466
Other Accession	P70052 , Q64104 , Q91379 , NP_003260.1
Reactivity	Human
Predicted	Chicken, Mouse, Xenopus
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB41747
Calculated MW	42589
Antigen Region	75-102

Additional Information

Gene ID	7101
Other Names	Nuclear receptor subfamily 2 group E member 1, Nuclear receptor TLX, Protein tailless homolog, TII, hTII, NR2E1, TLX
Target/Specificity	This NR2E1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 75-102 amino acids from the Central region of human NR2E1.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	NR2E1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	NR2E1
Synonyms	TLX

Function	Orphan receptor that binds DNA as a monomer to hormone response elements (HRE) containing an extended core motif half-site sequence 5'-AAGGTCA-3' in which the 5' flanking nucleotides participate in determining receptor specificity (By similarity). May be required to pattern anterior brain differentiation. Involved in the regulation of retinal development and essential for vision. During retinogenesis, regulates PTEN-Cyclin D expression via binding to the promoter region of PTEN and suppressing its activity (By similarity). May be involved in retinoic acid receptor (RAR) regulation in retinal cells.
Cellular Location	Nucleus {ECO:0000255 PROSITE-ProRule:PRU00407}.
Tissue Location	Brain specific. Present in all brain sections tested, highest levels in the caudate nucleus and hippocampus, weakest levels in the thalamus.


Background

Orphan receptor that binds DNA as a monomer to hormone response elements (HRE) containing an extended core motif half-site sequence 5'-AAGGTCA-3' in which the 5' flanking nucleotides participate in determining receptor specificity. May be required for brain development. May be involved in the regulation of retinal development (By similarity).

References

Liu, H.K., et al. Genes Dev. 24(7):683-695(2010)
Iwahara, N., et al. Biochem. Biophys. Res. Commun. 386(4):671-675(2009)
de Krom, M., et al. Biol. Psychiatry 65(7):625-630(2009)
Kumar, R.A., et al. Am. J. Med. Genet. B Neuropsychiatr. Genet. 147B (6), 880-889 (2008) :
Yokoyama, A., et al. Mol. Cell. Biol. 28(12):3995-4003(2008)

Images



NR2E1 Antibody (Center) (Cat. #AP19904c) western blot analysis in HL-60 cell line lysates (35ug/lane). This demonstrates the NR2E1 antibody detected the NR2E1 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.