10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# SCNN1B Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20003c

## Product Information

| | |
|---|---|
| **Application** | WB, E |
| **Primary Accession** | P51168 |
| **Other Accession** | NP_000327.2 |
| **Reactivity** | Human |
| **Predicted** | Mouse, Rat, Bovine, Canine, Rabbit |
| **Host** | Rabbit |
| **Clonality** | Polyclonal |
| **Isotype** | Rabbit IgG |
| **Clone Names** | RB42043 |
| **Calculated MW** | 72659 |
| **Antigen Region** | 306-332 |

## Additional Information

| | |
|---|---|
| **Gene ID** | 6338 |
| **Other Names** | Amiloride-sensitive sodium channel subunit beta, Beta-NaCH, Epithelial Na(+) channel subunit beta, Beta-ENaC, ENaCB, Nonvoltage-gated sodium channel 1 subunit beta, SCNEB, SCNN1B |
| **Target/Specificity** | This SCNN1B antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 306-332 amino acids from the Central region of human SCNN1B. |
| **Dilution** | WB~~1:1000 E~~Use at an assay dependent concentration. |
| **Format** | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| **Storage** | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| **Precautions** | SCNN1B Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures. |

## Protein Information

| | |
|---|---|
| **Name** | SCNN1B {ECO:0000303\|PubMed:7490094, ECO:0000312\|HGNC:HGNC:10600} |

| | |
|---|---|
| **Function** | This is one of the three pore-forming subunits of the heterotrimeric epithelial sodium channel (ENaC), a critical regulator of sodium balance and fluid homeostasis (PubMed:30251954, PubMed:32729833, PubMed:7762608, PubMed:9792722). ENaC operates in epithelial tissues, where it mediates the electrodiffusion of sodium ions from extracellular fluid through the apical membrane of cells, with water following osmotically (PubMed:24124190). It plays a key role in maintaining sodium homeostasis through electrogenic sodium reabsorption in the kidneys (PubMed:12107247). Additionally, ENaC is essential for airway surface liquid homeostasis, which is crucial for proper mucus clearance (PubMed:24124190). |
| **Cellular Location** | Apical cell membrane; Multi-pass membrane protein. Cytoplasmic vesicle membrane {ECO:0000250\|UniProtKB:P37090}; Multi-pass membrane protein |
| **Tissue Location** | Detected in placenta, lung and kidney (PubMed:7762608). Expressed in kidney (at protein level) (PubMed:22207244). |


## Background

Nonvoltage-gated, amiloride-sensitive, sodium channels control fluid and electrolyte transport across epithelia in many organs. These channels are heteromeric complexes consisting of 3 subunits: alpha, beta, and gamma. This gene encodes the beta subunit, and mutations in this gene have been associated with pseudohypoaldosteronism type 1 (PHA1), and Liddle syndrome.

## References


Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
Song, W., et al. J. Biol. Chem. 285(13):9716-9728(2010)
Yokoyama, K., et al. Nephron Clin Pract 115 (4), C237-C243 (2010) :
McGeachie, M., et al. Circulation 120(24):2448-2454(2009)
Azad, A.K., et al. Hum. Mutat. 30(7):1093-1103(2009)


## Images




SCNN1B Antibody (Center) (Cat. #AP20003c) western blot analysis in ZR-75-1 cell line lysates (35ug/lane).This demonstrates the SCNN1B antibody detected the SCNN1B protein (arrow).


Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.