

UBE2S Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20071b

Product Information

Application	WB, IHC-P, E
Primary Accession	Q16763
Other Accession	B5DFI8 , Q921J4 , Q1RML1 , A1L3K1 , NP_055316.2
Reactivity	Human
Predicted	Xenopus, Bovine, Mouse, Rat, Canine, Rabbit
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB42367
Calculated MW	23845
Antigen Region	183-210

Additional Information

Gene ID	27338
Other Names	Ubiquitin-conjugating enzyme E2 S, E2-EPF, Ubiquitin carrier protein S, Ubiquitin-conjugating enzyme E2-24 kDa, Ubiquitin-conjugating enzyme E2-EPF5, Ubiquitin-protein ligase S, UBE2S, E2EPF
Target/Specificity	This UBE2S antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 183-210 amino acids from the C-terminal region of human UBE2S.
Dilution	WB~~1:1000 IHC-P~~1:100 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	UBE2S Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	UBE2S
Synonyms	E2EPF

Function

Accepts ubiquitin from the E1 complex and catalyzes its covalent attachment to other proteins (PubMed:[19820702](#), PubMed:[19822757](#), PubMed:[22496338](#), PubMed:[27259151](#)). Catalyzes 'Lys-11'-linked polyubiquitination. Acts as an essential factor of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-regulated ubiquitin ligase that controls progression through mitosis (PubMed:[19820702](#), PubMed:[19822757](#), PubMed:[27259151](#), PubMed:[27910872](#)). Acts by specifically elongating 'Lys-11'-linked polyubiquitin chains initiated by the E2 enzyme UBE2C/UBCH10 on APC/C substrates, enhancing the degradation of APC/C substrates by the proteasome and promoting mitotic exit (PubMed:[19820702](#), PubMed:[19822757](#), PubMed:[27259151](#)). Also acts by elongating ubiquitin chains initiated by the E2 enzyme UBE2D1/UBCH5 in vitro; it is however unclear whether UBE2D1/UBCH5 acts as an E2 enzyme for the APC/C in vivo. Also involved in ubiquitination and subsequent degradation of VHL, resulting in an accumulation of HIF1A (PubMed:[16819549](#)). In vitro able to promote polyubiquitination using all 7 ubiquitin Lys residues, except 'Lys-48'-linked polyubiquitination (PubMed:[20061386](#), PubMed:[20622874](#)).


Background

This gene encodes a member of the ubiquitin-conjugating enzyme family. The encoded protein is able to form a thiol ester linkage with ubiquitin in a ubiquitin activating enzyme-dependent manner, a characteristic property of ubiquitin carrier proteins.

References


Wu, T., et al. Proc. Natl. Acad. Sci. U.S.A. 107(4):1355-1360(2010)
Garnett, M.J., et al. Nat. Cell Biol. 11(11):1363-1369(2009)
Williamson, A., et al. Proc. Natl. Acad. Sci. U.S.A. 106(43):18213-18218(2009)
Chen, M.F., et al. J. Mol. Med. 87(3):307-320(2009)
Lim, J.H., et al. J. Cell. Biochem. 105(4):1117-1127(2008)

Images



UBE2S Antibody (C-term) (Cat. #AP20071b) western blot analysis in HL-60 cell line lysates (35ug/lane). This demonstrates the UBE2S antibody detected the UBE2S protein (arrow).

Formalin-fixed and paraffin-embedded H.adrenal tissue reacted with UBE2S Antibody (C-term) (Cat#AP20071b).



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.