

SGOL2 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20106c

Product Information

Application	WB, E
Primary Accession	Q562F6
Other Accession	NP_689737.3
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	144739
Antigen Region	391-419

Additional Information

Gene ID	151246
Other Names	Shugoshin-like 2, Shugoshin-2, Sgo2, Tripin, SGOL2
Target/Specificity	This SGOL2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 391-419 amino acids from the Central region of human SGOL2.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	SGOL2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	SGO2 (HGNC:30812)
Synonyms	SGOL2
Function	Cooperates with PPP2CA to protect centromeric cohesin from separase-mediated cleavage in oocytes specifically during meiosis I. Has a crucial role in protecting REC8 at centromeres from cleavage by separase.

During meiosis, protects centromeric cohesion complexes until metaphase II/anaphase II transition, preventing premature release of meiosis-specific REC8 cohesin complexes from anaphase I centromeres. Is thus essential for an accurate gametogenesis. May act by targeting PPP2CA to centromeres, thus leading to cohesin dephosphorylation (By similarity). Essential for recruiting KIF2C to the inner centromere and for correcting defective kinetochore attachments. Involved in centromeric enrichment of AUKRB in prometaphase.

Cellular Location

Nucleus. Chromosome, centromere. Chromosome, centromere, kinetochore {ECO:0000250|UniProtKB:Q7TSY8}. Note=During meiosis I, accumulates at centromeres during diplotene, and colocalizes differentially with the cohesin subunits RAD21 and REC8 at metaphase I centromeres (By similarity). SGO2 and RAD21 change their relative distributions during telophase I when sister-kinetochore association is lost (By similarity). During meiosis II, it shows a striking tension- dependent redistribution within centromeres throughout chromosome congression during prometaphase II, as it does during mitosis (By similarity). In Hela cells, localizes at centromeres throughout prophase until metaphase and disappears at anaphase (PubMed:17485487) Centromeric localization requires the presence of BUB1 and AUKRB (PubMed:17485487). {ECO:0000250|UniProtKB:Q7TSY8, ECO:0000269|PubMed:17485487}


Background

Cooperates with PPP2CA to protect centromeric cohesin from separase-mediated cleavage in oocytes specifically during meiosis I. Has a crucial role in protecting REC8 at centromeres from cleavage by separase. During meiosis, protects centromeric cohesion complexes until metaphase II/anaphase II transition, preventing premature release of meiosis-specific REC8 cohesin complexes from anaphase I centromeres. Is thus essential for an accurate gametogenesis. May act by targeting PPP2CA to centromeres, thus leading to cohesin dephosphorylation (By similarity). Essential for recruiting KIF2C to the inner centromere and for correcting defective kinetochore attachments.

References

Tanno, Y., et al. Genes Dev. 24(19):2169-2179(2010)
Ross, C.J., et al. Nat. Genet. 41(12):1345-1349(2009)
Lee, J., et al. Nat. Cell Biol. 10(1):42-52(2008)
Huang, H., et al. J. Cell Biol. 177(3):413-424(2007)
Kitajima, T.S., et al. Nature 441(7089):46-52(2006)

Images



SGOL2 Antibody (Center) (Cat. #AP20106c) western blot analysis in U-937 cell line lysates (35ug/lane). This demonstrates the SGOL2 antibody detected the SGOL2 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.