

Mouse Txnip Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20130c

Product Information

Application	IHC-P, WB, E
Primary Accession	Q8BG60
Other Accession	Q5M7W1 , NP_001009935.1
Reactivity	Human, Rat, Mouse
Predicted	Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB30652
Calculated MW	44363
Antigen Region	116-145

Additional Information

Gene ID	56338
Other Names	Thioredoxin-interacting protein, Vitamin D3 up-regulated protein 1, Txnip, Vdup1
Target/Specificity	This Mouse Txnip antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 116-145 amino acids from the Central region of mouse Txnip.
Dilution	IHC-P~~1:100 WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Mouse Txnip Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	Txnip
Synonyms	Vdup1

Function	May act as an oxidative stress mediator by inhibiting thioredoxin activity or by limiting its bioavailability (PubMed: 10843682). Interacts with COPS5 and restores COPS5-induced suppression of CDKN1B stability, blocking the COPS5-mediated translocation of CDKN1B from the nucleus to the cytoplasm (PubMed: 15930262). Functions as a transcriptional repressor, possibly by acting as a bridge molecule between transcription factors and corepressor complexes, and over-expression will induce G0/G1 cell cycle arrest (By similarity). Required for the maturation of natural killer cells (PubMed: 15723808). Acts as a suppressor of tumor cell growth. Inhibits the proteasomal degradation of DDIT4, and thereby contributes to the inhibition of the mammalian target of rapamycin complex 1 (mTORC1) (By similarity).
Cellular Location	Cytoplasm.
Tissue Location	Ubiquitously expressed.

Background

May act as an oxidative stress mediator by inhibiting thioredoxin activity or by limiting its bioavailability. Interacts with COPS5 and restores COPS5-induced suppression of CDKN1B stability, blocking the COPS5-mediated translocation of CDKN1B from the nucleus to the cytoplasm. Functions as a transcriptional repressor, possibly by acting as a bridge molecule between transcription factors and corepressor complexes, and over-expression will induce G0/G1 cell cycle arrest. Required for the maturation of natural killer cells.

References


Kwon, H.J., et al. Toxicol. Appl. Pharmacol. 248(3):277-284(2010)
 Kwon, H.J., et al. J. Immunol. 185(7):3980-3989(2010)
 Ren, Y., et al. FEBS Lett. 584(15):3480-3485(2010)
 Chutkow, W.A., et al. Diabetes 59(6):1424-1434(2010)
 Shao, Y., et al. Immunol. Lett. 129(2):78-84(2010)

Images



Immunohistochemical analysis of AP20130c on paraffin-embedded Mouse liver tissue. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:100) for 1 hour at room temperature. Undiluted CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.

Anti-Mouse Txnip Antibody (Center) at 1:2000 dilution + Mouse stomach tissue lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 44 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.