

CTAGE1 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20180c

Product Information

Application	WB, E
Primary Accession	Q96RT6
Other Accession	NP_758441.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB41795
Calculated MW	85282
Antigen Region	386-414

Additional Information

Gene ID	64693
Other Names	cTAGE family member 2, Protein cTAGE-2, Cancer/testis antigen 212, CT212, CTAGE1, CTAGE2
Target/Specificity	This CTAGE1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 386-414 amino acids from the Central region of human CTAGE1.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	CTAGE1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CTAGE1
Synonyms	CTAGE2
Cellular Location	Membrane; Single-pass membrane protein

Tissue Location

Testis. Not found in tumor.

Background

The function of this protein is unknown.


References

Thye, T., et al. Nat. Genet. 42(9):739-741(2010)

Usener, D., et al. J. Invest. Dermatol. 121(1):198-206(2003)

Eichmuller, S., et al. Proc. Natl. Acad. Sci. U.S.A. 98(2):629-634(2001)

Images



Anti-CTAGE1 Antibody (Center) at 1:1000 dilution + human testis lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 85 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.