

DENR Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20311c

Product Information

Application	WB, E
Primary Accession	Q43583
Other Accession	Q9CQJ6 , Q2HJ47
Reactivity	Human, Mouse
Predicted	Bovine, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB40460
Calculated MW	22092
Antigen Region	47-76

Additional Information

Gene ID	8562
Other Names	Density-regulated protein, DRP, Protein DRP1, Smooth muscle cell-associated protein 3, SMAP-3, DENR, DRP1
Target/Specificity	This DENR antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 47-76 amino acids from the Central region of human DENR.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	DENR Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information


Name	DENR
Synonyms	DRP1

Function	Translation regulator forming a complex with MCTS1 to promote translation reinitiation. Translation reinitiation is the process where the small ribosomal subunit remains attached to the mRNA following termination of translation of a regulatory upstream ORF (uORF), and resume scanning on the same mRNA molecule to initiate translation of a downstream ORF, usually the main ORF (mORF). The MCTS1/DENR complex is pivotal to two linked mechanisms essential for translation reinitiation. Firstly, the dissociation of deacylated tRNAs from post- termination 40S ribosomal complexes during ribosome recycling. Secondly, the recruitment in an EIF2-independent manner of aminoacylated initiator tRNA to P site of 40S ribosomes for a new round of translation. This regulatory mechanism governs the translation of more than 150 genes which translation reinitiation is MCTS1/DENR complex-dependent.
Cellular Location	Cytoplasm.
Tissue Location	Highly expressed in heart and skeletal muscle and moderately expressed in the brain, placenta, liver and pancreas. Weakly expressed in the lung and kidney.

Background

May be involved in the translation of target mRNAs by scanning and recognition of the initiation codon. Plays a role in the modulation of the translational profile of a subset of cancer-related mRNAs when recruited to the translational initiation complex by the oncogene MCTS1.

Images



DENR Antibody (Center) (Cat. #AP20311c) western blot analysis in 293 cell line lysates (35ug/lane). This demonstrates the DENR antibody detected the DENR protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.