

# TP53 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20337a

## Product Information

---

|                          |  |
|--------------------------|--|
| <b>Application</b>       | WB, E  |
| <b>Primary Accession</b> | <a href="#">P04637</a>   |
| <b>Other Accession</b>   | <a href="#">Q95330</a> , <a href="#">Q9TUB2</a> , <a href="#">P56423</a> |
| <b>Reactivity</b>        | Human, Rat, Mouse  |
| <b>Predicted</b>         | Monkey, Pig, Rabbit  |
| <b>Host</b>              | Rabbit   |
| <b>Clonality</b>         | Polyclonal   |
| <b>Isotype</b>           | Rabbit IgG   |
| <b>Clone Names</b>       | RB42478  |
| <b>Calculated MW</b>     | 43653  |
| <b>Antigen Region</b>    | 1-30   |

## Additional Information

---

|                           |  |
|---------------------------|--|
| <b>Gene ID</b>            | 7157   |
| <b>Other Names</b>        | Cellular tumor antigen p53, Antigen NY-CO-13, Phosphoprotein p53, Tumor suppressor p53, TP53, P53  |
| <b>Target/Specificity</b> | This TP53 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human TP53.            |
| <b>Dilution</b>           | WB~~1:1000 E~~Use at an assay dependent concentration.   |
| <b>Format</b>             | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| <b>Storage</b>            | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.                                      |
| <b>Precautions</b>        | TP53 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.   |

## Protein Information

---

|                 |      |
|-----------------|------|
| <b>Name</b>     | TP53 |
| <b>Synonyms</b> | P53  |


|                          |   |
|--------------------------|---|
| <b>Function</b>          | <p>Multifunctional transcription factor that induces cell cycle arrest, DNA repair or apoptosis upon binding to its target DNA sequence (PubMed:<a href="#">11025664</a>, PubMed:<a href="#">12524540</a>, PubMed:<a href="#">12810724</a>, PubMed:<a href="#">15186775</a>, PubMed:<a href="#">15340061</a>, PubMed:<a href="#">17317671</a>, PubMed:<a href="#">17349958</a>, PubMed:<a href="#">19556538</a>, PubMed:<a href="#">20673990</a>, PubMed:<a href="#">20959462</a>, PubMed:<a href="#">22726440</a>, PubMed:<a href="#">24051492</a>, PubMed:<a href="#">24652652</a>, PubMed:<a href="#">35618207</a>, PubMed:<a href="#">36634798</a>, PubMed:<a href="#">38653238</a>, PubMed:<a href="#">9840937</a>). Acts as a tumor suppressor in many tumor types; induces growth arrest or apoptosis depending on the physiological circumstances and cell type (PubMed:<a href="#">11025664</a>, PubMed:<a href="#">12524540</a>, PubMed:<a href="#">12810724</a>, PubMed:<a href="#">15186775</a>, PubMed:<a href="#">15340061</a>, PubMed:<a href="#">17189187</a>, PubMed:<a href="#">17317671</a>, PubMed:<a href="#">17349958</a>, PubMed:<a href="#">19556538</a>, PubMed:<a href="#">20673990</a>, PubMed:<a href="#">20959462</a>, PubMed:<a href="#">22726440</a>, PubMed:<a href="#">24051492</a>, PubMed:<a href="#">24652652</a>, PubMed:<a href="#">38653238</a>, PubMed:<a href="#">9840937</a>). Negatively regulates cell division by controlling expression of a set of genes required for this process (PubMed:<a href="#">11025664</a>, PubMed:<a href="#">12524540</a>, PubMed:<a href="#">12810724</a>, PubMed:<a href="#">15186775</a>, PubMed:<a href="#">15340061</a>, PubMed:<a href="#">17317671</a>, PubMed:<a href="#">17349958</a>, PubMed:<a href="#">19556538</a>, PubMed:<a href="#">20673990</a>, PubMed:<a href="#">20959462</a>, PubMed:<a href="#">22726440</a>, PubMed:<a href="#">24051492</a>, PubMed:<a href="#">24652652</a>, PubMed:<a href="#">38653238</a>, PubMed:<a href="#">9840937</a>). One of the activated genes is an inhibitor of cyclin-dependent kinases. Apoptosis induction seems to be mediated either by stimulation of BAX and FAS antigen expression, or by repression of Bcl-2 expression (PubMed:<a href="#">12524540</a>, PubMed:<a href="#">17189187</a>). Its pro-apoptotic activity is activated via its interaction with PPP1R13B/ASPP1 or TP53BP2/ASPP2 (PubMed:<a href="#">12524540</a>). However, this activity is inhibited when the interaction with PPP1R13B/ASPP1 or TP53BP2/ASPP2 is displaced by PPP1R13L/iASPP (PubMed:<a href="#">12524540</a>). In cooperation with mitochondrial PPIF is involved in activating oxidative stress-induced necrosis; the function is largely independent of transcription. Induces the transcription of long intergenic non-coding RNA p21 (lincRNA-p21) and lincRNA-Mkln1. LincRNA-p21 participates in TP53-dependent transcriptional repression leading to apoptosis and seems to have an effect on cell-cycle regulation. Implicated in Notch signaling cross-over. Prevents CDK7 kinase activity when associated to CAK complex in response to DNA damage, thus stopping cell cycle progression. Isoform 2 enhances the transactivation activity of isoform 1 from some but not all TP53-inducible promoters. Isoform 4 suppresses transactivation activity and impairs growth suppression mediated by isoform 1. Isoform 7 inhibits isoform 1-mediated apoptosis. Regulates the circadian clock by repressing CLOCK-BMAL1-mediated transcriptional activation of PER2 (PubMed:<a href="#">24051492</a>).</p> |
| <b>Cellular Location</b> | <p>Cytoplasm. Nucleus. Nucleus, PML body. Endoplasmic reticulum. Mitochondrion matrix. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome Note=Recruited into PML bodies together with CHEK2 (PubMed:<a href="#">12810724</a>) Translocates to mitochondria upon oxidative stress (PubMed:<a href="#">22726440</a>) Translocates to mitochondria in response to mitomycin C treatment (PubMed:<a href="#">27323408</a>). Competitive inhibition of TP53 interaction with HSPA9/MOT-2 by UBXN2A results in increased protein abundance and subsequent translocation of TP53 to the nucleus (PubMed:<a href="#">24625977</a>) [Isoform 2]: Nucleus. Cytoplasm. Note=Localized mainly in the nucleus with minor staining in the cytoplasm [Isoform 4]: Nucleus. Cytoplasm. Note=Predominantly nuclear but translocates to the cytoplasm following cell stress [Isoform 8]: Nucleus. Cytoplasm. Note=Localized in both nucleus and cytoplasm in most cells. In some cells, forms foci in the nucleus that are different from nucleoli</p>   |
| <b>Tissue Location</b>   | <p>Ubiquitous. Isoforms are expressed in a wide range of normal tissues but in a tissue-dependent manner. Isoform 2 is expressed in most normal tissues but is not detected in brain, lung, prostate, muscle, fetal brain, spinal cord and</p>  |

fetal liver. Isoform 3 is expressed in most normal tissues but is not detected in lung, spleen, testis, fetal brain, spinal cord and fetal liver. Isoform 7 is expressed in most normal tissues but is not detected in prostate, uterus, skeletal muscle and breast. Isoform 8 is detected only in colon, bone marrow, testis, fetal brain and intestine. Isoform 9 is expressed in most normal tissues but is not detected in brain, heart, lung, fetal liver, salivary gland, breast or intestine

## Background

Acts as a tumor suppressor in many tumor types; induces growth arrest or apoptosis depending on the physiological circumstances and cell type. Involved in cell cycle regulation as a trans-activator that acts to negatively regulate cell division by controlling a set of genes required for this process. One of the activated genes is an inhibitor of cyclin-dependent kinases. Apoptosis induction seems to be mediated either by stimulation of BAX and FAS antigen expression, or by repression of Bcl-2 expression. Implicated in Notch signaling cross-over. Prevents CDK7 kinase activity when associated to CAK complex in response to DNA damage, thus stopping cell cycle progression. Isoform 2 enhances the transactivation activity of isoform 1 from some but not all TP53-inducible promoters. Isoform 4 suppresses transactivation activity and impairs growth suppression mediated by isoform 1. Isoform 7 inhibits isoform 1-mediated apoptosis.

## Images



All lanes : Anti-TP53 Antibody (N-term) at 1:2000 dilution  
Lane 1: HeLa whole cell lysate Lane 2: MCF-7 whole cell lysate Lane 3: NIH/3T3 whole cell lysate Lane 4: PC-12 whole cell lysate Lysates/proteins at 20 µg per lane.  
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 44 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.