

CPVL Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP20351b

Product Information

Application WB, E **Primary Accession Q9H3G5** Reactivity Human Host Rabbit Clonality Polyclonal Isotype Rabbit IgG **Clone Names** RB42793 **Calculated MW** 54164 **Antigen Region** 448-476

Additional Information

Gene ID 54504

Other Names Probable serine carboxypeptidase CPVL, 3416-, Carboxypeptidase,

vitellogenic-like, Vitellogenic carboxypeptidase-like protein, VCP-like protein,

hVLP, CPVL, VLP

Target/Specificity This CPVL antibody is generated from rabbits immunized with a KLH

conjugated synthetic peptide between 448-476 amino acids from the

C-terminal region of human CPVL.

Dilution WB~~1:1000 E~~Use at an assay dependent concentration.

Format Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions CPVL Antibody (C-term) is for research use only and not for use in diagnostic

or therapeutic procedures.

Protein Information

Name CPVL

Synonyms VLP

Function May be involved in the digestion of phagocytosed particles in the lysosome,

participation in an inflammatory protease cascade, and trimming of peptides for antigen presentation.


Tissue Location

Expressed in macrophages but not in other leukocytes. Abundantly expressed in heart and kidney. Also expressed in spleen, leukocytes, and placenta

Background

May be involved in the digestion of phagocytosed particles in the lysosome, participation in an inflammatory protease cascade, and trimming of peptides for antigen presentation.

Images



All lanes: Anti-CPVL Antibody (C-term) at 1:1000 dilution Lane 1: 293 whole cell lysate Lane 2: HepG2 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 54 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.