

ETV7 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20366b

Product Information

Application	WB, E
Primary Accession	Q9Y603
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB42837
Calculated MW	38998
Antigen Region	313-341

Additional Information

Gene ID	51513
Other Names	Transcription factor ETV7, ETS translocation variant 7, ETS-related protein Tel2, Tel-related Ets factor, Transcription factor Tel-2, ETV7, TEL2, TELB, TREF
Target/Specificity	This ETV7 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 313-341 amino acids from the C-terminal region of human ETV7.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ETV7 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ETV7
Synonyms	TEL2, TELB, TREF
Function	Transcriptional repressor; binds to the DNA sequence 5'- CCGGAAGT-3'. Isoform A does not seem to have a repressor activity. Isoform C does not

seem to have a repressor activity.

Cellular Location

Nucleus.


Tissue Location

Expressed in hematopoietic tissues.

Background

Transcriptional repressor; binds to the DNA sequence 5'-CCGGAAGT-3'. Isoform A does not seem to have a repressor activity. Isoform C does not seem to have a repressor activity.

Images



All lanes : Anti-ETV7 Antibody (C-term) at 1:2000 dilution
Lane 1: A431 whole cell lysate Lane 2: Jurkat whole cell lysate Lane 3: K562 whole cell lysate Lane 4: NCI-H292 whole cell lysate Lane 5: U-87 MG whole cell lysate
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 39 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.