10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

FBXO28 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20438b

Product Information

Application	WB, E
Primary Accession	Q9NVF7
Other Accession	Q8BIG4, Q2NL16
Reactivity	Human, Mouse
Predicted	Bovine, Rat, Canine, Rabbit
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	41149
Antigen Region	339-368

Additional Information

Gene ID	23219
Other Names	F-box only protein 28, FBXO28, KIAA0483
Target/Specificity	This FBXO28 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 339-368 amino acids from the C-terminal region of human FBXO28.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	FBXO28 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	FBXO28
Synonyms	CENP-30 {ECO:0000303|PubMed:20813266}, K
Function	Substrate-recognition component of the SCF (SKP1-CUL1-F-box protein)-type E3 ubiquitin ligase complex, promoting ubiquitination and proteasomal

	degradation of specific target proteins including TOP2A, RAB27A or itself (PubMed:27754753, PubMed:31678254). Regulates topoisomerase IIalpha/TOP2A decatenation activity and plays an important role in maintaining genomic stability (PubMed:27754753). Plays a role in lipid metabolism and inflammation through the ubiquitinated degradation of RAB27A (By similarity). Strongly regulates beta-cell survival without having any significant independent effect on insulin secretion (PubMed:29587369). Plays an essential role in spindle morphology and actin-based spindle migration probably through the ARPC2/ARP3 signaling pathway (By similarity).
Cellular Location	Chromosome, centromere, kinetochore. Nucleus Chromosome {ECO:0000250|UniProtKB:Q8BIG4}. Note=Localizes in the vicinity of mitotic chromosomes during mitosis


Background

Probably recognizes and binds to some phosphorylated proteins and promotes their ubiquitination and degradation (By similarity).

References

Ota T., et al. Nat. Genet. 36:40-45(2004).
Gregory S.G., et al. Nature 441:315-321(2006).
Seki N., et al. DNA Res. 4:345-349(1997).
Olsen J.V., et al. Cell 127:635-648(2006).
Dephoure N., et al. Proc. Natl. Acad. Sci. U.S.A. 105:10762-10767(2008).

Images



FBXO28 Antibody (C-term) (Cat. #AP20438b) western blot analysis in Hela,MCF-7,ZR-75-1 cell line and mouse brain lysates (35ug/lane).This demonstrates the FBXO28 antibody detected the FBXO28 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.