

TRIM6 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP20469a

Product Information

Application	WB, E
Primary Accession	<u>Q9C030</u>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	56400
Antigen Region	26-55

Additional Information

Gene ID	117854
Other Names	Tripartite motif-containing protein 6, RING finger protein 89, TRIM6, RNF89
Target/Specificity	This TRIM6 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 26-55 amino acids from the N-terminal region of human TRIM6.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	TRIM6 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	TRIM6
Synonyms	RNF89
Function	E3 ubiquitin ligase that plays a crucial role in the activation of the IKBKE-dependent branch of the type I interferon signaling pathway (PubMed: <u>24882218</u> , PubMed: <u>31694946</u>). In concert with the ubiquitin-conjugating E2 enzyme UBE2K, synthesizes unanchored 'Lys-

48'-linked polyubiquitin chains that promote the oligomerization and autophosphorylation of IKBKE leading to stimulation of an antiviral response (PubMed:<u>24882218</u>). Also ubiquitinates MYC and inhibits its transcription activation activity, maintaining the pluripotency of embryonic stem cells (By similarity). Promotes the association of unanchored 'Lys-48'-polyubiquitin chains with DHX16 leading to enhanced RIGI-mediated innate antiviral immune response (PubMed:<u>35263596</u>).


Cellular Location

Cytoplasm

References

Reymond A., et al. EMBO J. 20:2140-2151(2001). Ota T., et al. Nat. Genet. 36:40-45(2004). Mural R.J., et al. Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases. Orimo A., et al. Genomics 69:143-149(2000).

Images



All lanes : Anti-TRIM6 Antibody (N-term) at 1:2000 dilution Lane 1: Ramos whole cell lysate Lane 2: U-2OS whole cell lysate Lane 3: PC-3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 56 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.