

M Cckar Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20509b

Product Information

Application	WB, E
Primary Accession	O08786
Other Accession	P30551
Reactivity	Mouse
Predicted	Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	48437
Antigen Region	325-356

Additional Information

Gene ID	12425
Other Names	Cholecystokinin receptor type A, CCK-A receptor, CCK-AR, Cholecystokinin-1 receptor, CCK1-R, Cckar
Target/Specificity	This Mouse Cckar antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 325-356 amino acids from the C-terminal region of mouse Cckar.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	M Cckar Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	Cckar
Function	Receptor for cholecystokinin. Mediates pancreatic growth and enzyme secretion, smooth muscle contraction of the gall bladder and stomach. Has a 1000-fold higher affinity for CCK rather than for gastrin. It modulates feeding

and dopamine-induced behavior in the central and peripheral nervous system. This receptor mediates its action by association with G proteins that activate a phosphatidylinositol-calcium second messenger system (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein.

Background

Receptor for cholecystikinin. Mediates pancreatic growth and enzyme secretion, smooth muscle contraction of the gall bladder and stomach. Has a 1000-fold higher affinity for CCK rather than for gastrin. It modulates feeding and dopamine-induced behavior in the central and peripheral nervous system. This receptor mediates its action by association with G proteins that activate a phosphatidylinositol-calcium second messenger system (By similarity).


References

Takata Y., et al. Gene 187:267-271(1997).

Lacourse K.A., et al. Biochem. Biophys. Res. Commun. 236:630-635(1997).

Carninci P., et al. Science 309:1559-1563(2005).

Images



Mouse Cckar Antibody (C-term) (Cat. #AP20509b) western blot analysis in mouse bladder, cerebellum and lung tissue lysates (35ug/lane). This demonstrates the Mouse Cckar antibody detected the Mouse Cckar protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.