10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

BLMH Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20513c

Product Information

Application	WB, IHC-P, E
Primary Accession	Q13867
Other Accession	P70645, P13019, Q8R016
Reactivity	Human, Rat, Mouse
Predicted	Rabbit, Bovine, Canine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	52562
Antigen Region	212-242

Additional Information

Gene ID	642
Other Names	Bleomycin hydrolase, BH, BLM hydrolase, BMH, BLMH
Target/Specificity	This BLMH antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 212-242 amino acids from the Central region of human BLMH.
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	BLMH Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	BLMH
Function	The normal physiological role of BLM hydrolase is unknown, but it catalyzes the inactivation of the antitumor drug BLM (a glycopeptide) by hydrolyzing the carboxamide bond of its B- aminoalaninamide moiety thus protecting normal and malignant cells from BLM toxicity.

Cellular Location	Cytoplasm. Cytoplasmic granule. Note=Co-localizes with NUDT12 in the cytoplasmic granules.


Background

The normal physiological role of BLM hydrolase is unknown, but it catalyzes the inactivation of the antitumor drug BLM (a glycopeptide) by hydrolyzing the carboxamide bond of its B-aminoalaninamide moiety thus protecting normal and malignant cells from BLM toxicity (By similarity).


References

Barrow I.K.-P., et al. Submitted (AUG-1998) to the EMBL/GenBank/DDBJ databases.
Ferrando A.A., et al. Cancer Res. 56:1746-1750(1996).
Broemme D., et al. Biochemistry 35:6706-6714(1996).
Kalnine N., et al. Submitted (OCT-2004) to the EMBL/GenBank/DDBJ databases.
Ota T., et al. Nat. Genet. 36:40-45(2004).

Images



All lanes : Anti-BLMH Antibody (Center) at 1:1000 dilution Lane 1: HL-60 whole cell lysate Lane 2: MOLT-4 whole cell lysate Lane 3: Neuro-2a whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 53 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Immunohistochemical analysis of paraffin-embedded H. skin section using BLMH Antibody (Center)(Cat#AP20513C). AP20513C was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.