10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

R Cdk4 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20515b

Product Information

Application	WB, IF, IHC-P, E
Primary Accession	P35426
Reactivity	Human, Rat, Mouse
Predicted	Rabbit
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	33799
Antigen Region	272-303

Additional Information

Gene ID	94201
Other Names	Cyclin-dependent kinase 4, Cell division protein kinase 4, PSK-J3, Cdk4
Target/Specificity	This Rat Cdk4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 272-303 amino acids from the C-terminal region of rat Cdk4.
Dilution	WB~~1:1000 IF~~1:25 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	R Cdk4 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	Cdk4
Function	Ser/Thr-kinase component of cyclin D-CDK4 (DC) complexes that phosphorylate and inhibit members of the retinoblastoma (RB) protein family including RB1 and regulate the cell-cycle during G(1)/S transition. Phosphorylation of RB1 allows dissociation of the transcription factor E2F

	from the RB/E2F complexes and the subsequent transcription of E2F target genes which are responsible for the progression through the G(1) phase. Hypophosphorylates RB1 in early G(1) phase. Cyclin D-CDK4 complexes are major integrators of various mitogenenic and antimitogenic signals. Also phosphorylates SMAD3 in a cell-cycle-dependent manner and represses its transcriptional activity. Component of the ternary complex, cyclin D/CDK4/CDKN1B, required for nuclear translocation and activity of the cyclin D-CDK4 complex (By similarity).
Cellular Location	Cytoplasm {ECO:0000250|UniProtKB:P11802}. Nucleus {ECO:0000250|UniProtKB:P11802}. Nucleus membrane {ECO:0000250|UniProtKB:P11802}. Note=Cytoplasmic when non-complexed Forms a cyclin D-CDK4 complex in the cytoplasm as cells progress through G(1) phase. The complex accumulates on the nuclear membrane and enters the nucleus on transition from G(1) to S phase. Also present in nucleoli and heterochromatin lumps. Colocalizes with RB1 after release into the nucleus (By similarity). {ECO:0000250|UniProtKB:P11802}
Tissue Location	Expressed in fetal and adult lung. Also expressed in brain, heart, liver, skeletal muscle and testes


Background

Ser/Thr-kinase component of cyclin D-CDK4 (DC) complexes that phosphorylate and inhibit members of the retinoblastoma (RB) protein family including RB1 and regulate the cell-cycle during G(1)/S transition. Phosphorylation of RB1 allows dissociation of the transcription factor E2F from the RB/E2F complexes and the subsequent transcription of E2F target genes which are responsible for the progression through the G(1) phase. Hypophosphorylates RB1 in early G(1) phase. Cyclin D-CDK4 complexes are major integrators of various mitogenenic and antimitogenic signals. Also phosphorylates SMAD3 in a cell-cycle-dependent manner and represses its transcriptional activity. Component of the ternary complex, cyclin D/CDK4/CDKN1B, required for nuclear translocation and activity of the cyclin D-CDK4 complex (By similarity).

References


Cho F.S., et al. Biochem. Biophys. Res. Commun. 191:860-865(1993).

Images




Immunohistochemical analysis of paraffin-embedded R. brain section using R Cdk4 Antibody (C-term)(Cat#AP20515B). AP20515B was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.


Immunohistochemical analysis of paraffin-embedded rat small intestine section using Rat Cdk4 Antibody (C-term)(Cat#AP20515B). AP20515B was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB



staining.



Fluorescent image of MCF-7 cells stained with (Rat) Cdk4 Antibody (C-term)(Cat#AP20515B). AP20515B was diluted at 1:25 dilution. An Alexa Fluor 488-conjugated goat anti-rabbit lgG at 1:400 dilution was used as the secondary antibody (green). Cytoplasmic actin was counterstained with Alexa Fluor® 555 conjugated with Phalloidin (red).



Rat Cdk4 Antibody (C-term) (Cat. #AP20515b) western blot analysis in rat lung tissue lysates (35ug/lane).This demonstrates the rat Cdk4 antibody detected the rat Cdk4 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.