10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

KHDRBS2 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20541b

Product Information

Application	WB, E
Primary Accession	Q5VWX1
Reactivity	Mouse, Rat, Human
Predicted	Bovine, Canine, Rabbit
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	38927
Antigen Region	298-321

Additional Information

Gene ID	202559
Other Names	KH domain-containing, RNA-binding, signal transduction-associated protein 2, Sam68-like mammalian protein 1, SLM-1, hSLM-1, KHDRBS2, SLM1
Target/Specificity	This KHDRBS2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 298-321 amino acids from the C-terminal region of human KHDRBS2.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	KHDRBS2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	KHDRBS2
Synonyms	SLM1
Function	RNA-binding protein that plays a role in the regulation of alternative splicing and influences mRNA splice site selection and exon inclusion. Binds both

poly(A) and poly(U) homopolymers. Phosphorylation by PTK6 inhibits its RNA-binding ability (By similarity). Induces an increased concentration-dependent incorporation of exon in CD44 pre- mRNA by direct binding to purine-rich exonic enhancer. Can regulate alternative splicing of NRXN1 in the laminin G-like domain 6 containing the evolutionary conserved neurexin alternative spliced segment 4 (AS4) involved in neurexin selective targeting to postsynaptic partners. Regulates cell-type specific alternative splicing of NRXN1 at AS4 and acts synergystically with SAM68 in exon skipping. In contrast acts antagonistically with SAM68 in NRXN3 exon skipping at AS4. Its phosphorylation by FYN inhibits its ability to regulate splice site selection. May function as an adapter protein for Src kinases during mitosis.

Cellular Location Nucleus {ECO:0000250|UniProtKB:Q9WU01}.

Tissue Location Highly expressed in brain, lung, kidney and small intestine. Weakly expressed in placenta, liver, spleen, thymus, ovary and colon.


Background

RNA-binding protein that plays a role in the regulation of alternative splicing and influences mRNA splice site selection and exon inclusion. Its phosphorylation by FYN inhibits its ability to regulate splice site selection. Induces an increased concentration-dependent incorporation of exon in CD44 pre-mRNA by direct binding to purine-rich exonic enhancer. May function as an adapter protein for Src kinases during mitosis. Binds both poly(A) and poly(U) homopolymers. Phosphorylation by PTK6 inhibits its RNA-binding ability (By similarity).

References

Wang L., et al. Mol. Biol. Rep. 29:369-375(2002).
Ota T., et al. Nat. Genet. 36:40-45(2004).
Mungall A.J., et al. Nature 425:805-811(2003).
Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Cote J., et al. Mol. Biol. Cell 14:274-287(2003).

Images



KHDRBS2 Antibody (C-term) (Cat. #AP20541b) western blot analysis in Hela cell line and rat testis tissue lysates (35ug/lane).This demonstrates the KHDRBS2 antibody detected the KHDRBS2 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.