

# NDUFA9 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20542c

## Product Information

---

<b>Application</b>	WB, E
<b>Primary Accession</b>	<a href="#">Q16795</a>
<b>Reactivity</b>	Mouse, Rat, Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Calculated MW</b>	42510
<b>Antigen Region</b>	99-121

## Additional Information

---

<b>Gene ID</b>	4704
<b>Other Names</b>	NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 9, mitochondrial, Complex I-39kD, CI-39kD, NADH-ubiquinone oxidoreductase 39 kDa subunit, NDUFA9, NDUF52L
<b>Target/Specificity</b>	This NDUFA9 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 99-121 amino acids from the Central region of human NDUFA9.
<b>Dilution</b>	WB~~1:1000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	NDUFA9 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	NDUFA9
<b>Synonyms</b>	NDUF52L
<b>Function</b>	Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed not to be involved in catalysis.

Required for proper complex I assembly (PubMed:[28671271](#)). Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.

#### Cellular Location

Mitochondrion matrix

## Background

---

Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed not to be involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.


## References

---

- Baens M., et al. Genomics 16:214-218(1993).  
Loeffen J.L.C.M., et al. Submitted (FEB-1998) to the EMBL/GenBank/DDBJ databases.  
Cross S.H., et al. Nat. Genet. 6:236-244(1994).  
Murray J., et al. J. Biol. Chem. 278:13619-13622(2003).  
Burkard T.R., et al. BMC Syst. Biol. 5:17-17(2011).

## Images

---



NDUFA9 Antibody (Center) (Cat. #AP20542c) western blot analysis in HeLa, Jurkat cell line and mouse heart tissue lysates (35ug/lane). This demonstrates the NDUFA9 antibody detected the NDUFA9 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.