

JAM3 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20552b

Product Information

Application	WB, E
Primary Accession	Q9BX67
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB47643
Calculated MW	35020

Additional Information

Gene ID	83700
Other Names	Junctional adhesion molecule C, JAM-C, JAM-2, Junctional adhesion molecule 3, JAM-3, JAM3
Target/Specificity	This JAM3 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 261-295 amino acids from the C-terminal region of human JAM3.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	JAM3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	JAM3
Function	Junctional adhesion protein that mediates heterotypic cell- cell interactions with its cognate receptor JAM2 to regulate different cellular processes (PubMed: 11590146 , PubMed: 11823489). Plays a role in homing and mobilization of hematopoietic stem and progenitor cells within the bone marrow. At the surface of bone marrow stromal cells, it contributes to the

retention of the hematopoietic stem and progenitor cells expressing JAM3 (PubMed:[11590146](#), PubMed:[24357068](#)). Plays a central role in leukocytes extravasation by facilitating transmigration through the endothelium (By similarity). Plays a role in spermatogenesis where JAM2 and JAM3, which are respectively expressed by Sertoli and germ cells, mediate an interaction between both cell types and play an essential role in the anchorage of germ cells onto Sertoli cells and the assembly of cell polarity complexes during spermatid differentiation (By similarity). Also functions as a counter- receptor for ITGAM, mediating leukocyte-platelet interactions and is involved in the regulation of transepithelial migration of polymorphonuclear neutrophils (PMN) (PubMed:[12208882](#), PubMed:[15194813](#)). Plays a role in angiogenesis (PubMed:[23255084](#)). Plays a role in the regulation of cell migration (Probable). During myogenesis, it is involved in myocyte fusion (By similarity).

Cellular Location

Cell membrane; Single-pass type I membrane protein. Cell junction. Cell junction, desmosome. Cell junction, tight junction. Note=Detected in the acrosome region in developing spermatids (By similarity). In epithelial cells, it is expressed at desmosomes but not at tight junctions (PubMed:15194813) Localizes at the cell surface of endothelial cells; treatment of endothelial cells with vascular endothelial growth factor stimulates recruitment of JAM3 to cell-cell contacts (PubMed:15994945) {ECO:0000250|UniProtKB:Q9D8B7}

Tissue Location

Detected on round and elongated spermatids (at protein level) (PubMed:15372036). Highest expression in placenta, brain and kidney. Significant expression is detected on platelets. Expressed in intestinal mucosa cells. Expressed in the vascular endothelium Found in serum (at protein level). Also detected in the synovial fluid of patients with rheumatoid arthritis, psoriatic arthritis or osteoarthritis (at protein level).

Background


Participates in cell-cell adhesion. It is a counter- receptor for ITGAM, mediating leukocyte-platelet interactions and is involved in the regulation of transepithelial migration of polymorphonuclear neutrophils (PMN). The soluble form is a mediator of angiogenesis.

References

Arrate M.P.,et al.J. Biol. Chem. 276:45826-45832(2001).
Aurrand-Lions M.A.,et al.Blood 98:3699-3707(2001).
Santoso S.,et al.J. Exp. Med. 196:679-691(2002).
Phillips H.M.,et al.Genomics 79:475-478(2002).
Ota T.,et al.Nat. Genet. 36:40-45(2004).

Images

All lanes: Anti-JAM3 Antibody (C-term) at 1:500 dilution +
Mouse brain lysate Lysates/proteins at 20 µg per lane.
Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase
conjugated (ASP1615) at 1/15000 dilution. Observed band
size: 35 KDa Blocking/Dilution buffer: 5% NFDM/TBST.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.