10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

ANP32E Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20559a

Product Information

Application	WB, IF, E
Primary Accession	Q9BTT0
Other Accession	Q5XIE0, P97822
Reactivity	Human, Rat, Mouse
Predicted	Rat, Bovine, Canine, Rabbit, Chicken
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB44588
Calculated MW	30692

Additional Information

Gene ID	81611
Other Names	Acidic leucine-rich nuclear phosphoprotein 32 family member E, LANP-like protein, LANP-L, ANP32E
Target/Specificity	This ANP32E antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 21-44 amino acids from the N-terminal region of human ANP32E.
Dilution	WB~~1:1000 IF~~1:25 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ANP32E Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ANP32E
Function	Histone chaperone that specifically mediates the genome-wide removal of histone H2A.Z/H2AZ1 from the nucleosome: removes H2A.Z/H2AZ1 from its normal sites of deposition, especially from enhancer and insulator regions.

	Not involved in deposition of H2A.Z/H2AZ1 in the nucleosome. May stabilize the evicted H2A.Z/H2AZ1-H2B dimer, thus shifting the equilibrium towards dissociation and the off-chromatin state (PubMed:24463511). Inhibits activity of protein phosphatase 2A (PP2A). Does not inhibit protein phosphatase 1. May play a role in cerebellar development and synaptogenesis.
Cellular Location	Cytoplasm. Nucleus.
Tissue Location	Expressed in peripheral blood leukocytes, colon, small intestine, prostate, thymus, spleen, skeletal muscle, liver and kidney.


Background

Inhibits activity of protein phosphatase 2A (PP2A). Does not inhibit protein phosphatase 1. May play a role in cerebellar development and synaptogenesis process by modulating PP2A activity (By similarity).


References

Jiang M.,et al.Cytogenet. Genome Res. 97:68-71(2002). Ota T.,et al.Nat. Genet. 36:40-45(2004). Bechtel S.,et al.BMC Genomics 8:399-399(2007). Gregory S.G.,et al.Nature 441:315-321(2006). Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.

Images



Fluorescent image of SH-SY5Y cells stained with ANP32E Antibody (N-term)(Cat#AP20559a). AP20559a was diluted at 1:25 dilution. An Alexa Fluor 488-conjugated goat anti-rabbit lgG at 1:400 dilution was used as the secondary antibody (green). Cytoplasmic actin was counterstained with Alexa Fluor® 555 conjugated with Phalloidin (red).



Western blot analysis of lysates from SH-SY5Y, U-937 cell line and mouse brain tissue lysate (from left to right), using ANP32E Antibody (N-term) (Cat. #AP20559a). AP20559a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug per lane.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.