

AK4 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20571a

Product Information

Application	WB, IHC-P, IF, E
Primary Accession	P27144
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB48826
Calculated MW	25268

Additional Information

Gene ID	205
Other Names	Adenylate kinase 4, mitochondrial {ECO:0000255 HAMAP-Rule:MF_03170}, AK4 {ECO:0000255 HAMAP-Rule:MF_03170}, 27410 {ECO:0000255 HAMAP-Rule:MF_03170}, 2746 {ECO:0000255 HAMAP-Rule:MF_03170}, Adenylate kinase 3-like {ECO:0000255 HAMAP-Rule:MF_03170}, GTP:AMP phosphotransferase AK4 {ECO:0000255 HAMAP-Rule:MF_03170}, AK4 {ECO:0000255 HAMAP-Rule:MF_03170}
Target/Specificity	This AK4 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 119-153 amino acids from the Central region of human AK4.
Dilution	WB~~1:1000 IHC-P~~1:100~500 IF~~1:25 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	AK4 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	AK4 (HGNC:363)
-------------	----------------------------------

Function	Broad-specificity mitochondrial nucleoside phosphate kinase involved in cellular nucleotide homeostasis by catalyzing nucleoside- phosphate interconversions (PubMed: 19073142 , PubMed: 19766732 , PubMed: 23416111 , PubMed: 24767988). Similar to other adenylate kinases, preferentially catalyzes the phosphorylation of the nucleoside monophosphate AMP with ATP as phosphate donor to produce ADP (PubMed: 19766732). Phosphorylates only AMP when using GTP as phosphate donor (PubMed: 19766732). In vitro, can also catalyze the phosphorylation of CMP, dAMP and dCMP and use GTP as an alternate phosphate donor (PubMed: 19766732 , PubMed: 23416111). Moreover, exhibits a diphosphate kinase activity, producing ATP, CTP, GTP, UTP, TTP, dATP, dCTP and dGTP from the corresponding diphosphate substrates with either ATP or GTP as phosphate donors (PubMed: 23416111). Plays a role in controlling cellular ATP levels by regulating phosphorylation and activation of the energy sensor protein kinase AMPK (PubMed: 24767988 , PubMed: 26980435). Plays a protective role in the cellular response to oxidative stress (PubMed: 19130895 , PubMed: 23474458 , PubMed: 26980435).
Cellular Location	Mitochondrion matrix {ECO:0000255 HAMAP- Rule:MF_03170, ECO:0000269 PubMed:11485571, ECO:0000269 PubMed:19766732, ECO:0000269 PubMed:26980435}
Tissue Location	Highly expressed in kidney, moderately expressed in heart and liver and weakly expressed in brain


Background

Involved in maintaining the homeostasis of cellular nucleotides by catalyzing the interconversion of nucleoside phosphates. Efficiently phosphorylates AMP and dAMP using ATP as phosphate donor, but phosphorylates only AMP when using GTP as phosphate donor. Also displays broad nucleoside diphosphate kinase activity.

References


Xu G.,et al.Genomics 13:537-542(1992).
 Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.
 Ota T.,et al.Nat. Genet. 36:40-45(2004).
 Gregory S.G.,et al.Nature 441:315-321(2006).
 Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.

Images



All lanes : Anti-AK4 Antibody (Center)at 1:1000 dilution +
 Raji whole cell lysate Lysates/proteins at 20 μ g per lane.
 Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase
 conjugated (ASP1615) at 1/15000 dilution. Observed band
 size : 26kDa Blocking/Dilution buffer: 5% NFDM/TBST.

All lanes : Anti-AK4 Antibody (Center)at 1:1000 dilution



Lane 1:Mouse kidney lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size : 26kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Citations

- [MiR-199a-3p affects the multi-chemoresistance of osteosarcoma through targeting AK4.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.