10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

NDUFC2 Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP20601A

Product Information

Application	WB, FC, IF, IHC-P, E
Primary Accession	O95298
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB48893
Calculated MW	14188

Additional Information

Gene ID	4718
Other Names	NADH dehydrogenase [ubiquinone] 1 subunit C2, Complex I-B145b, CI-B145b, Human lung cancer oncogene 1 protein, HLC-1, NADH-ubiquinone oxidoreductase subunit B145b, NDUFC2
Target/Specificity	This NDUFC2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 5-39 amino acids from the N-terminal region of human NDUFC2.
Dilution	WB~~1:1000 FC~~1:25 IF~~1:25 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	NDUFC2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	NDUFC2 (HGNC:7706)
Function	Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed not to be involved in catalysis but required for the complex assembly. Complex I functions in the transfer of

	electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.
Cellular Location	Mitochondrion inner membrane; Single-pass membrane protein; Matrix side


Background

Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed not to be involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.


References

Loeffen J.L.C.M.,et al.Biochem. Biophys. Res. Commun. 253:415-422(1998).
Dai F.Y.,et al.Submitted (AUG-1998) to the EMBL/GenBank/DDBJ databases.
Zhang Q.-H.,et al.Genome Res. 10:1546-1560(2000).
Wiemann S.,et al.Genome Res. 11:422-435(2001).
Kim J.W.,et al.Submitted (APR-2001) to the EMBL/GenBank/DDBJ databases.

Images





All lanes: Anti-NDUFC2 Antibody (N-term) at 1:1000 dilution Lane 1: 293T whole cell lysate Lane 2: Hela whole cell lysate Lane 3: Jurkat whole cell lysate Lane 4: Mouse brain lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 11 KDa Blocking/Dilution buffer: 5% NFDM/TBST.



Flow cytometric analysis of HepG2 cells using NDUFC2 Antibody (N-term)(green, Cat#AP20601a) compared to an isotype control of rabbit IgG(blue). AP20601a was diluted at 1:25 dilution. An Alexa Fluor® 488 goat anti-rabbit lgG at 1:400 dilution was used as the secondary antibody.

Fluorescent image of HepG2 cells stained with NDUFC2 Antibody (N-term) (Cat#AP20601a). AP20601a was diluted at 1:25 dilution. An Alexa Fluor 488-conjugated goat anti-rabbit lgG at 1:400 dilution was used as the secondary antibody (green). DAPI was used to stain the cell nuclear (blue).





Immunohistochemical analysis of paraffin-embedded H. kidney section using NDUFC2 Antibody (N-term)(Cat#AP20601a). AP20601a was diluted at 1:100 dilution. A peroxidase-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.