

NDUFC2 Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20601A

Product Information

| | |
|--------------------------|------------------------|
| Application | WB, FC, IF, IHC-P, E |
| Primary Accession | O95298 |
| Reactivity | Human, Mouse, Rat |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | Rabbit IgG |
| Clone Names | RB48893 |
| Calculated MW | 14188 |

Additional Information

| | |
|---------------------------|---|
| Gene ID | 4718 |
| Other Names | NADH dehydrogenase [ubiquinone] 1 subunit C2, Complex I-B145b, CI-B145b, Human lung cancer oncogene 1 protein, HLC-1, NADH-ubiquinone oxidoreductase subunit B145b, NDUFC2 |
| Target/Specificity | This NDUFC2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 5-39 amino acids from the N-terminal region of human NDUFC2. |
| Dilution | WB~~1:1000 FC~~1:25 IF~~1:25 IHC-P~~1:100~500 E~~Use at an assay dependent concentration. |
| Format | Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| Storage | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| Precautions | NDUFC2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures. |

Protein Information

| | |
|-----------------|--|
| Name | NDUFC2 (HGNC:7706) |
| Function | Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed not to be involved in catalysis but required for the complex assembly. Complex I functions in the transfer of |

electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.

Cellular Location

Mitochondrion inner membrane; Single-pass membrane protein; Matrix side


Background

Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed not to be involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.


References

Loeffen J.L.C.M.,et al.Biochem. Biophys. Res. Commun. 253:415-422(1998).
Dai F.Y.,et al.Submitted (AUG-1998) to the EMBL/GenBank/DDBJ databases.
Zhang Q.-H.,et al.Genome Res. 10:1546-1560(2000).
Wiemann S.,et al.Genome Res. 11:422-435(2001).
Kim J.W.,et al.Submitted (APR-2001) to the EMBL/GenBank/DDBJ databases.

Images





All lanes: Anti-NDUFC2 Antibody (N-term) at 1:1000 dilution Lane 1: 293T whole cell lysate Lane 2: HeLa whole cell lysate Lane 3: Jurkat whole cell lysate Lane 4: Mouse brain lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 11 KDa Blocking/Dilution buffer: 5% NFDm/TBST.



Flow cytometric analysis of HepG2 cells using NDUFC2 Antibody (N-term)(green, Cat#AP20601a) compared to an isotype control of rabbit IgG(blue). AP20601a was diluted at 1:25 dilution. An Alexa Fluor® 488 goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody.

Fluorescent image of HepG2 cells stained with NDUFC2 Antibody (N-term) (Cat#AP20601a). AP20601a was diluted at 1:25 dilution. An Alexa Fluor 488-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody (green). DAPI was used to stain the cell nuclear (blue).



Immunohistochemical analysis of paraffin-embedded H. kidney section using NDUFC2 Antibody (N-term)(Cat#AP20601a). AP20601a was diluted at 1:100 dilution. A peroxidase-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.