

GSTT1 Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP20614a

Product Information

Application WB, E **Primary Accession** P30711 Reactivity Human, Rat Host Rabbit Clonality Polyclonal Isotype Rabbit IgG **Clone Names** RB47963 **Calculated MW** 27335

Additional Information

Gene ID 2952

Other Names Glutathione S-transferase theta-1, GST class-theta-1, Glutathione transferase

T1-1, GSTT1

Target/Specificity This GSTT1 antibody is generated from a rabbit immunized with a KLH

conjugated synthetic peptide between 46-79 amino acids from the N-terminal

region of human GSTT1.

Dilution WB~~1:1000 E~~Use at an assay dependent concentration.

Format Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein A column, followed by peptide

affinity purification.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions GSTT1 Antibody (N-term) is for research use only and not for use in diagnostic

or therapeutic procedures.

Protein Information

Name GSTT1

Function Conjugation of reduced glutathione to a wide number of exogenous and

endogenous hydrophobic electrophiles. Acts on 1,2-epoxy-

3-(4-nitrophenoxy)propane, phenethylisothiocyanate 4-nitrobenzyl chloride and 4-nitrophenethyl bromide. Displays glutathione peroxidase activity with

cumene hydroperoxide.

Cellular Location Cytoplasm.

Tissue Location Found in erythrocyte. Expressed at low levels in liver. In lung, expressed at

low levels in club cells and ciliated cells at the alveolar/bronchiolar junction.

Absent from epithelial cells of larger bronchioles.


Background

Conjugation of reduced glutathione to a wide number of exogenous and endogenous hydrophobic electrophiles. Acts on 1,2- epoxy-3-(4-nitrophenoxy)propane, phenethylisothiocyanate 4- nitrobenzyl chloride and 4-nitrophenethyl bromide. Displays glutathione peroxidase activity with cumene hydroperoxide.

References

Pemble S.,et al.Biochem. J. 300:271-276(1994). Jemth P.,et al.Arch. Biochem. Biophys. 348:247-254(1997). Sprenger R.,et al.Pharmacogenetics 10:557-565(2000). Iida A.,et al.Submitted (MAR-2001) to the EMBL/GenBank/DDBJ databases. Kalnine N.,et al.Submitted (OCT-2004) to the EMBL/GenBank/DDBJ databases.

Images



Western blot analysis of lysates from human brain, rat liver and lung tissue lysate (from left to right), using GSTT1 Antibody (N-term)(Cat. #AP20614a). AP20614a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.