

HTR1A Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20616c

Product Information

Application	WB, FC, E
Primary Accession	P08908
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB46792
Calculated MW	46107

Additional Information

Gene ID	3350
Other Names	5-hydroxytryptamine receptor 1A, 5-HT-1A, 5-HT1A, G-21, Serotonin receptor 1A, HTR1A, ADRB2RL1, ADRBRL1
Target/Specificity	This HTR1A antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 239-273 amino acids from the Central region of human HTR1A.
Dilution	WB~~1:1000 FC~~1:25 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.05% (V/V) Proclin 300. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	HTR1A Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	HTR1A (HGNC:5286)
Synonyms	ADRB2RL1, ADRBRL1
Function	G-protein coupled receptor for 5-hydroxytryptamine (serotonin) (PubMed: 22957663 , PubMed: 3138543 , PubMed: 33762731 , PubMed: 37935376 , PubMed: 37935377 , PubMed: 8138923 , PubMed: 8393041).

Also functions as a receptor for various drugs and psychoactive substances (PubMed:[22957663](#), PubMed:[3138543](#), PubMed:[33762731](#), PubMed:[38552625](#), PubMed:[8138923](#), PubMed:[8393041](#)). Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of downstream effectors, such as adenylate cyclase (PubMed:[22957663](#), PubMed:[3138543](#), PubMed:[33762731](#), PubMed:[8138923](#), PubMed:[8393041](#)). HTR1A is coupled to G(i)/G(o) G alpha proteins and mediates inhibitory neurotransmission: signaling inhibits adenylate cyclase activity and activates a phosphatidylinositol-calcium second messenger system that regulates the release of Ca(2+) ions from intracellular stores (PubMed:[33762731](#), PubMed:[35610220](#)). Beta-arrestin family members regulate signaling by mediating both receptor desensitization and resensitization processes (PubMed:[18476671](#), PubMed:[20363322](#), PubMed:[20945968](#)). Plays a role in the regulation of 5- hydroxytryptamine release and in the regulation of dopamine and 5- hydroxytryptamine metabolism (PubMed:[18476671](#), PubMed:[20363322](#), PubMed:[20945968](#)). Plays a role in the regulation of dopamine and 5- hydroxytryptamine levels in the brain, and thereby affects neural activity, mood and behavior (PubMed:[18476671](#), PubMed:[20363322](#), PubMed:[20945968](#)). Plays a role in the response to anxiogenic stimuli (PubMed:[18476671](#), PubMed:[20363322](#), PubMed:[20945968](#)).

Cellular Location

Cell membrane; Multi-pass membrane protein. Cell projection, dendrite {ECO:0000250|UniProtKB:P19327}

Tissue Location

Detected in lymph nodes, thymus and spleen. Detected in activated T-cells, but not in resting T-cells

Background


G-protein coupled receptor for 5-hydroxytryptamine (serotonin). Also functions as a receptor for various drugs and psychoactive substances. Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors, such as adenylate cyclase. Beta-arrestin family members inhibit signaling via G proteins and mediate activation of alternative signaling pathways. Signaling inhibits adenylate cyclase activity and activates a phosphatidylinositol-calcium second messenger system that regulates the release of Ca(2+) ions from intracellular stores. Plays a role in the regulation of 5- hydroxytryptamine release and in the regulation of dopamine and 5- hydroxytryptamine metabolism. Plays a role in the regulation of dopamine and 5-hydroxytryptamine levels in the brain, and thereby affects neural activity, mood and behavior. Plays a role in the response to anxiogenic stimuli.

References


- Kobilka B.K.,et al.Nature 329:75-79(1987).
 Saltzman A.G.,et al.Submitted (FEB-1991) to the EMBL/GenBank/DDBJ databases.
 Levy F.O.,et al.Submitted (MAY-1992) to the EMBL/GenBank/DDBJ databases.
 Kitano T.,et al.Mol. Biol. Evol. 21:936-944(2004).
 Puhl H.L. III,et al.Submitted (APR-2002) to the EMBL/GenBank/DDBJ databases.

Images

All lanes : Anti-HTR1A Antibody (Center) at 1:1000 dilution
 Lane 1: Molt-4 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Observed



band size : 46kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Flow cytometric analysis of Jurkat cells using HTR1A Antibody (Center)(green, Cat#AP20616c) compared to an isotype control of rabbit IgG(blue). AP20616c was diluted at 1:25 dilution. An Alexa Fluor® 488 goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.