

DLG2 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20636c

Product Information

Application	WB, E
Primary Accession	Q15700
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB47835
Calculated MW	97552

Additional Information

Gene ID	1740
Other Names	Disks large homolog 2, Channel-associated protein of synapse-110, Chapsyn-110, Postsynaptic density protein PSD-93, DLG2
Target/Specificity	This DLG2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 614-648 amino acids from the Central region of human DLG2.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	DLG2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	DLG2
Function	Required for perception of chronic pain through NMDA receptor signaling. Regulates surface expression of NMDA receptors in dorsal horn neurons of the spinal cord. Interacts with the cytoplasmic tail of NMDA receptor subunits as well as inward rectifying potassium channels. Involved in regulation of synaptic stability at cholinergic synapses. Part of the postsynaptic protein

scaffold of excitatory synapses (By similarity).

Cellular Location

Cell membrane {ECO:0000250|UniProtKB:Q63622}; Lipid-anchor {ECO:0000250|UniProtKB:Q63622}. Postsynaptic density {ECO:0000250|UniProtKB:Q63622}. Synapse. Membrane {ECO:0000250|UniProtKB:Q63622}. Cell projection, axon {ECO:0000250|UniProtKB:Q63622}. Perikaryon {ECO:0000250|UniProtKB:Q63622}. Note=Concentrated in soma and postsynaptic density of a subset of neurons {ECO:0000250|UniProtKB:Q63622}


Background

Required for perception of chronic pain through NMDA receptor signaling. Regulates surface expression of NMDA receptors in dorsal horn neurons of the spinal cord. Interacts with the cytoplasmic tail of NMDA receptor subunits as well as inward rectifying potassium channels. Involved in regulation of synaptic stability at cholinergic synapses. Part of the postsynaptic protein scaffold of excitatory synapses (By similarity).

References

Kim E.,et al.Neuron 17:103-113(1996).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Bechtel S.,et al.BMC Genomics 8:399-399(2007).
Taylor T.D.,et al.Nature 440:497-500(2006).
Totoki Y.,et al.Submitted (MAR-2005) to the EMBL/GenBank/DBJ databases.

Images



Western blot analysis of tissue lysates from mouse brain and rat brain (from left to right), using DLG2 Antibody (Center)(Cat. #AP20636c). AP20636c was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.