

# GDF5 Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP2067a

## Product Information

---

<b>Application</b>	WB, IHC-P, E
<b>Primary Accession</b>	<a href="#">P43026</a>
<b>Other Accession</b>	<a href="#">NP_000548</a>
<b>Reactivity</b>	Human
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	Rabbit IgG
<b>Calculated MW</b>	55395
<b>Antigen Region</b>	13-41

## Additional Information

---

<b>Gene ID</b>	8200
<b>Other Names</b>	Growth/differentiation factor 5, GDF-5, Bone morphogenetic protein 14, BMP-14, Cartilage-derived morphogenetic protein 1, CDMP-1, Lipopolysaccharide-associated protein 4, LAP-4, LPS-associated protein 4, Radotermis, GDF5, BMP14, CDMP1
<b>Target/Specificity</b>	This GDF5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 13-41 amino acids from the N-terminal region of human GDF5.
<b>Dilution</b>	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	GDF5 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	GDF5
<b>Synonyms</b>	BMP14, CDMP1

<b>Function</b>	Growth factor involved in bone and cartilage formation. During cartilage development regulates differentiation of chondrogenic tissue through two pathways. Firstly, positively regulates differentiation of chondrogenic tissue through its binding of high affinity with BMPR1B and of less affinity with BMPR1A, leading to induction of SMAD1-SMAD5-SMAD8 complex phosphorylation and then SMAD protein signaling transduction (PubMed: <a href="#">15530414</a> , PubMed: <a href="#">21976273</a> , PubMed: <a href="#">24098149</a> , PubMed: <a href="#">25092592</a> ). Secondly, negatively regulates chondrogenic differentiation through its interaction with NOG (PubMed: <a href="#">21976273</a> ). Required to prevent excessive muscle loss upon denervation. This function requires SMAD4 and is mediated by phosphorylated SMAD1/5/8 (By similarity). Binds bacterial lipopolysaccharide (LPS) and mediates LPS-induced inflammatory response, including TNF secretion by monocytes (PubMed: <a href="#">11276205</a> ).
<b>Cellular Location</b>	Secreted. Cell membrane
<b>Tissue Location</b>	Predominantly expressed in long bones during embryonic development. Expressed in monocytes (at protein level)


## Background

GDF5 is a member of the bone morphogenetic protein (BMP) family and the TGF-beta superfamily. This group of proteins is characterized by a polybasic proteolytic processing site which is cleaved to produce a mature protein containing seven conserved cysteine residues. The members of this family are regulators of cell growth and differentiation in both embryonic and adult tissues. Mutations in this gene are associated with acromesomelic dysplasia, Hunter-Thompson type; brachydactyly, type C; and chondrodysplasia, Grebe type. These associations confirm that the gene product plays a role in skeletal development.

## References


Kusafuka, K., et al., *Virchows Arch.* 442(5):482-490 (2003).  
 Faiyaz-UI-Haque, M., et al., *Am. J. Med. Genet.* 111(1):31-37 (2002).  
 Everman, D.B., et al., *Am. J. Med. Genet.* 112(3):291-296 (2002).  
 Faiyaz-UI-Haque, M., et al., *Clin. Genet.* 61(6):454-458 (2002).  
 Ducy, P., et al., *Kidney Int.* 57(6):2207-2214 (2000).

## Images



The anti-GDF5 N-term Pab (Cat. #AP2067a) is used in Western blot to detect GDF5 in A549 cell lysate.

Formalin-fixed and paraffin-embedded human lung carcinoma tissue reacted with GDF5 antibody (N-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data



demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

## Citations

---

- [GDF-5 is suppressed by IL-1beta and enhances TGF-beta3-mediated chondrogenic differentiation in human rheumatoid fibroblast-like synoviocytes.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.