

AFAP1-Y451 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20715a

Product Information

Application	WB, E
Primary Accession	Q8N556
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB45037
Calculated MW	80725

Additional Information

Gene ID	60312
Other Names	Actin filament-associated protein 1, 110 kDa actin filament-associated protein, AFAP-110, AFAP1, AFAP
Target/Specificity	This antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 444-477 amino acids from human.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	AFAP1-Y451 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	AFAP1
Synonyms	AFAP
Function	Can cross-link actin filaments into both network and bundle structures (By similarity). May modulate changes in actin filament integrity and induce lamellipodia formation. May function as an adapter molecule that links other proteins, such as SRC and PKC to the actin cytoskeleton. Seems to play a role

in the development and progression of prostate adenocarcinoma by regulating cell-matrix adhesions and migration in the cancer cells.

Cellular Location

Cytoplasm, cytoskeleton, stress fiber

Tissue Location

Low expression in normal breast epithelial cell line MCF-10A and in tumorigenic breast cancer cell lines MCF-7, T-47D and ZR-75-1. Highly expressed in the invasive breast cancer cell lines MDA-MB-231 and MDA-MB-435. Overexpressed in prostate carcinoma


Background

Can cross-link actin filaments into both network and bundle structures (By similarity). May modulate changes in actin filament integrity and induce lamellipodia formation. May function as an adapter molecule that links other proteins, such as SRC and PKC to the actin cytoskeleton. Seems to play a role in the development and progression of prostate adenocarcinoma by regulating cell-matrix adhesions and migration in the cancer cells.

References

Han B.,et al.J. Biol. Chem. 279:54793-54801(2004).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Hillier L.W.,et al.Nature 434:724-731(2005).
Totoki Y.,et al.Submitted (MAR-2005) to the EMBL/GenBank/DBJ databases.
Olsen J.V.,et al.Cell 127:635-648(2006).

Images



Western blot analysis of lysates from MCF-7, MDA-MB-453 cell line (from left to right), using PAFAP1-Y451(Cat. #AP20715a). AP20715a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.