

MT-ND1 Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20750a

Product Information

| | |
|--------------------------|------------------------|
| Application | WB, E |
| Primary Accession | P03886 |
| Reactivity | Human |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | Rabbit IgG |
| Clone Names | RB50705 |
| Calculated MW | 35661 |

Additional Information

| | |
|---------------------------|--|
| Gene ID | 4535 |
| Other Names | NADH-ubiquinone oxidoreductase chain 1, NADH dehydrogenase subunit 1, MT-ND1, MTND1, NADH1, ND1 |
| Target/Specificity | This MT-ND1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 34-68 amino acids from the N-terminal region of human MT-ND1. |
| Dilution | WB~~1:1000 E~~Use at an assay dependent concentration. |
| Format | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| Storage | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| Precautions | MT-ND1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures. |

Protein Information

| | |
|-----------------|---|
| Name | MT-ND1 |
| Synonyms | MTND1, NADH1, ND1 |
| Function | Core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) which catalyzes electron transfer from NADH through the respiratory chain, using ubiquinone as an electron acceptor |

(PubMed:[1959619](#)). Essential for the catalytic activity and assembly of complex I (PubMed:[1959619](#), PubMed:[26929434](#)).

Cellular Location

Mitochondrion inner membrane {ECO:0000250 | UniProtKB:P03887};
Multi-pass membrane protein


Background

Core subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I) that is believed to belong to the minimal assembly required for catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone (By similarity).

References

Anderson S.,et al.Nature 290:457-465(1981).
Horai S.,et al.Proc. Natl. Acad. Sci. U.S.A. 92:532-536(1995).
Moilanen J.S.,et al.Mol. Biol. Evol. 20:2132-2142(2003).
Ingman M.,et al.Nature 408:708-713(2000).
Ingman M.,et al.Genome Res. 13:1600-1606(2003).

Images



Western blot analysis of lysates from HeLa, HepG2 cell line (from left to right), using MT-ND1 Antibody (N-term)(Cat. #AP20750a). AP20750a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.