

GDF15 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20785c

Product Information

Application	WB, E
Primary Accession	Q99988
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB48204
Calculated MW	34140

Additional Information

Gene ID	9518
Other Names	Growth/differentiation factor 15, GDF-15, Macrophage inhibitory cytokine 1, MIC-1, NSAID-activated gene 1 protein, NAG-1, NSAID-regulated gene 1 protein, NRG-1, Placental TGF-beta, Placental bone morphogenetic protein, Prostate differentiation factor, GDF15, MIC1, PDF, PLAB, PTGFB
Target/Specificity	This GDF15 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 258-293 amino acids from the C-terminal region of human GDF15.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	GDF15 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	GDF15 {ECO:0000303 PubMed:23468844, ECO:0000312 HGNC:HGNC:30142}
Function	Hormone produced in response to various stresses to confer information about those stresses to the brain, and trigger an aversive response,

characterized by nausea, vomiting, and/or loss of appetite (PubMed:[23468844](#), PubMed:[24971956](#), PubMed:[28846097](#), PubMed:[28846098](#), PubMed:[28846099](#), PubMed:[28953886](#), PubMed:[29046435](#), PubMed:[30639358](#), PubMed:[31875646](#), PubMed:[33589633](#), PubMed:[38092039](#)). The aversive response is both required to reduce continuing exposure to those stresses at the time of exposure and to promote avoidance behavior in the future (PubMed:[30639358](#), PubMed:[33589633](#), PubMed:[38092039](#)). Acts by binding to its receptor, GFRAL, activating GFRAL-expressing neurons localized in the area postrema and nucleus tractus solitarius of the brainstem (PubMed:[28846097](#), PubMed:[28846098](#), PubMed:[28846099](#), PubMed:[28953886](#), PubMed:[31535977](#)). It then triggers the activation of neurons localized within the parabrachial nucleus and central amygdala, which constitutes part of the 'emergency circuit' that shapes responses to stressful conditions (PubMed:[28953886](#)). The GDF15-GFRAL signal induces expression of genes involved in metabolism, such as lipid metabolism in adipose tissues (PubMed:[31402172](#)). Required for avoidance behavior in response to food allergens: induced downstream of mast cell activation to promote aversion and minimize harmful effects of exposure to noxious substances (By similarity). In addition to suppress appetite, also promotes weight loss by enhancing energy expenditure in muscle: acts by increasing calcium futile cycling in muscle (By similarity). Contributes to the effect of metformin, an anti-diabetic drug, on appetite reduction and weight loss: produced in the kidney in response to metformin treatment, thereby activating the GDF15-GFRAL response, leading to reduced appetite and weight (PubMed:[31875646](#), PubMed:[37060902](#)). The contribution of GDF15 to weight loss following metformin treatment is however limited and subject to discussion (PubMed:[36001956](#)). Produced in response to anticancer drugs, such as camptothecin or cisplatin, promoting nausea, vomiting and contributing to malnutrition (By similarity). Overproduced in many cancers, promoting anorexia in cancer (cachexia) (PubMed:[32661391](#)). Responsible for the risk of nausea and vomiting during pregnancy: high levels of GDF15 during pregnancy, mostly originating from the fetus, are associated with increased nausea and vomiting (PubMed:[38092039](#)). Maternal sensitivity to nausea is probably determined by pre-pregnancy exposure to GDF15, women with naturally high level of GDF15 being less susceptible to nausea than women with low levels of GDF15 before pregnancy (PubMed:[38092039](#)). Promotes metabolic adaptation in response to systemic inflammation caused by bacterial and viral infections in order to promote tissue tolerance and prevent tissue damage (PubMed:[31402172](#)). Required for tissue tolerance in response to myocardial infarction by acting as an inhibitor of leukocyte integrating activation, thereby protecting against cardiac rupture (By similarity). Inhibits growth hormone signaling on hepatocytes (By similarity).

Cellular Location

Secreted Note=Secreted in the plasma.


Tissue Location

Detected in plasma (at protein level) (PubMed:[28572090](#), PubMed:[29046435](#)). Highly expressed in placenta, with lower levels in prostate and colon and some expression in kidney (PubMed:[37060902](#), PubMed:[9348093](#)).

References

- Hromas R.,et al.Biochim. Biophys. Acta 1354:40-44(1997).
 Yokoyama-Kobayashi M.,et al.J. Biochem. 122:622-626(1997).
 Bootcov M.R.,et al.Proc. Natl. Acad. Sci. U.S.A. 94:11514-11519(1997).
 Paralkar V.M.,et al.J. Biol. Chem. 273:13760-13767(1998).
 Kalnine N.,et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.

Images



Western blot analysis of lysate from HT-1080 cell line, using GDF15 Antibody (C-term)(Cat. #AP20785c). AP20785c was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.