

Mouse Csk Antibody (N-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20789a

Product Information

Application	WB, IHC-P, E
Primary Accession	P41241
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB50568

Additional Information

Other Names	Tyrosine-protein kinase CSK, C-Src kinase, Protein-tyrosine kinase MPK-2, p50CSK, Csk
Target/Specificity	This Mouse Csk antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 67-101 amino acids from the N-terminal region of human Mouse Csk.
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	Mouse Csk Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Background


Non-receptor tyrosine-protein kinase that plays an important role in the regulation of cell growth, differentiation, migration and immune response. Phosphorylates tyrosine residues located in the C-terminal tails of Src-family kinases (SFKs) including LCK, SRC, HCK, FYN, LYN or YES1. Upon tail phosphorylation, Src-family members engage in intramolecular interactions between the phosphotyrosine tail and the SH2 domain that result in an inactive conformation. To inhibit SFKs, CSK is recruited to the plasma membrane via binding to transmembrane proteins or adapter proteins located near the plasma membrane. Suppresses signaling by various surface receptors, including T- cell receptor (TCR) and B-cell receptor (BCR) by

phosphorylating and maintaining inactive several positive effectors such as FYN or LCK (By similarity).


References

Klages S.,et al.Proc. Natl. Acad. Sci. U.S.A. 91:2597-2601(1994).
Farber C.R.,et al.Submitted (JAN-2005) to the EMBL/GenBank/DDBJ databases.
Carninci P.,et al.Science 309:1559-1563(2005).
Gibaldi-Hebenstreit P.,et al.Oncogene 7:2499-2506(1992).
Lubec G.,et al.Submitted (JAN-2009) to UniProtKB.

Images



Immunohistochemical analysis of paraffin-embedded M.spleen section using Mouse Csk Antibody (N-term)(Cat#AP20789a). AP20789a was diluted at 1:25 dilution. A peroxidase-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.



Western blot analysis of lysate from mouse liver tissue lysate, using Mouse Csk Antibody (N-term)(Cat.#AP20789a). AP20789a was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.