

HIST1H3B3-S10 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20798b

Product Information

Application	WB, E
Primary Accession	P68431
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB46416
Calculated MW	15404

Additional Information

Gene ID	8350;8351;8352;8353;8354;8355;8356;8357;8358;8968
Other Names	Histone H31, Histone H3/a, Histone H3/b, Histone H3/c, Histone H3/d, Histone H3/f, Histone H3/h, Histone H3/i, Histone H3/j, Histone H3/k, Histone H3/l, HIST1H3A, H3FA
Target/Specificity	This antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 8~39 amino acids from human.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	HIST1H3B3-S10 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	H3C1 (HGNC:4766)
Synonyms	H3FA, HIST1H3A
Function	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription

regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Cellular Location

Nucleus. Chromosome.


Background

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

References

Zhong R.,et al.Nucleic Acids Res. 11:7409-7425(1983).
Marashi F.,et al.Biochem. Cell Biol. 64:277-289(1986).
Albig W.,et al.Genomics 10:940-948(1991).
Kardalidou E.,et al.J. Cell. Biochem. 52:375-383(1993).
Runge D.,et al.Submitted (OCT-1994) to the EMBL/GenBank/DBJ databases.

Images



Western blot analysis of lysate from mouse C2C12 cell line, using Phospho-HIST1H3B3-S10. ctrl(Cat. #AP20798b). AP20798b was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 35ug.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.