

CTSD(heavy chain) Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20803c

Product Information

Application	WB, E
Primary Accession	P07339
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB50732
Calculated MW	44552

Additional Information

Gene ID	1509
Other Names	Cathepsin D, Cathepsin D light chain, Cathepsin D heavy chain, CTSD, CPSD
Target/Specificity	This CTSD(heavy chain) antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 241-273 amino acids from the Central region of human CTSD(heavy chain).
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	CTSD(heavy chain) Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CTSD
Synonyms	CPSD
Function	Acid protease active in intracellular protein breakdown. Plays a role in APP processing following cleavage and activation by ADAM30 which leads to APP degradation (PubMed: 27333034). Involved in the pathogenesis of several diseases such as breast cancer and possibly Alzheimer disease.

Cellular Location	Lysosome. Melanosome. Secreted, extracellular space. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV. In aortic samples, detected as an extracellular protein loosely bound to the matrix (PubMed:20551380)
Tissue Location	Expressed in the aorta extracellular space (at protein level) (PubMed:20551380). Expressed in liver (at protein level) (PubMed:1426530).


Background

Acid protease active in intracellular protein breakdown. Involved in the pathogenesis of several diseases such as breast cancer and possibly Alzheimer disease.

References

Faust P.L.,et al.Proc. Natl. Acad. Sci. U.S.A. 82:4910-4914(1985).
Westley B.R.,et al.Nucleic Acids Res. 15:3773-3786(1987).
Redecker B.,et al.DNA Cell Biol. 10:423-431(1991).
Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.
Kalnine N.,et al.Submitted (OCT-2004) to the EMBL/GenBank/DDBJ databases.

Images



Western blot analysis of lysates from A431, MCF-7, SK-BR-3 cell line (from left to right), using CTSD (heavy chain) Antibody (Center) (Cat. #AP20803c). AP20803c was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L (HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 35ug per lane.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.