

# CD19 Antibody (C-term)

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP20806c

## Product Information

---

Application	WB, E
Primary Accession	<a href="#">P15391</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB50733
Calculated MW	61128

## Additional Information

---

Gene ID	930
Other Names	B-lymphocyte antigen CD19, B-lymphocyte surface antigen B4, Differentiation antigen CD19, T-cell surface antigen Leu-12, CD19, CD19
Target/Specificity	This CD19 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 517-551 amino acids from the C-terminal region of human CD19.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	CD19 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	CD19
Function	Functions as a coreceptor for the B-cell antigen receptor complex (BCR) on B-lymphocytes (PubMed: <a href="#">29523808</a> ). Decreases the threshold for activation of downstream signaling pathways and for triggering B-cell responses to antigens (PubMed: <a href="#">1373518</a> , PubMed: <a href="#">16672701</a> , PubMed: <a href="#">2463100</a> ). Activates signaling pathways that lead to the activation of phosphatidylinositol 3-kinase

and the mobilization of intracellular Ca(2+) stores (PubMed:[12387743](#), PubMed:[16672701](#), PubMed:[9317126](#), PubMed:[9382888](#)). Is not required for early steps during B cell differentiation in the blood marrow (PubMed:[9317126](#)). Required for normal differentiation of B-1 cells (By similarity). Required for normal B cell differentiation and proliferation in response to antigen challenges (PubMed:[1373518](#), PubMed:[2463100](#)). Required for normal levels of serum immunoglobulins, and for production of high-affinity antibodies in response to antigen challenge (PubMed:[12387743](#), PubMed:[16672701](#), PubMed:[9317126](#)).

#### Cellular Location

Cell membrane; Single-pass type I membrane protein. Membrane raft {ECO:0000250|UniProtKB:P25918}; Single-pass type I membrane protein {ECO:0000250|UniProtKB:P25918}

#### Tissue Location

Detected on marginal zone and germinal center B cells in lymph nodes (PubMed:2463100). Detected on blood B cells (at protein level) (PubMed:16672701, PubMed:2463100)


## Background

Assembles with the antigen receptor of B-lymphocytes in order to decrease the threshold for antigen receptor-dependent stimulation.

## References


Stamenkovic I.,et al.J. Exp. Med. 168:1205-1210(1988).  
 Tedder T.F.,et al.J. Immunol. 143:712-717(1989).  
 Kozmik Z.,et al.Mol. Cell. Biol. 12:2662-2672(1992).  
 Zhou L.J.,et al.Immunogenetics 35:102-111(1992).  
 Kuroki K.,et al.Genes Immun. 3:S21-S30(2002).

## Images



Western blot analysis of lysates from Daudi, Raji cell line (from left to right), using CD19 Antibody (C-term)(Cat. #AP20806c). AP20806c was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 35ug per lane.

Western blot analysis of lysate from Ramos cell line, using CD19 Antibody (C-term)(Cat. #AP20806c). AP20806c was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 35ug.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.